

## FACTSHEET



# PATIENTS AT RISK OF LYMPHOEDEMA WHO NEED SURGERY

Please note that this factsheet is being discontinued. Information for patients on lymphoedema is available on the Macmillan Cancer Support website (<u>https://bit.ly/Macmillan-lymphoedema</u>) and the Lymphoedema Support Network (<u>lymphoedema.org</u>).

Lymphoedema is a condition in which lymphatic fluid does not drain properly from an arm, leg or breast, causing swelling and discomfort. You can read more at the Macmillan cancer information and support website here: bit.ly/3s2qNx6

Every patient who has had cancer surgery involving the removal of the glands (lymph nodes) can be at risk of developing lymphoedema in their limbs, although the risk is small. It can also uncommonly develop following other types of surgery. Radiotherapy, chemotherapy and obesity can also increase the risk.

Lymphoedema may develop several years after surgery, which is why it is important to tell doctors and nurses about any previous cancer surgery you have had. It can also be associated with weight gain.

If you have an operation under general anaesthesia, healthcare staff at the hospital will use a blood pressure cuff to measure your blood pressure and will put a cannula (a thin plastic tube) into a vein in your arm. If you are having a hand or wrist operation, the surgeon usually uses a tourniquet (a tight band) during surgery to reduce blood loss.

If you have lymphoedema, these procedures could make lymphoedema worse. If you are having such a procedure, whenever possible, staff will use the arm on the other side. Sometimes they may mark the 'at risk' arm to remind everyone not to use it. If they have to use a tourniquet on that limb, they will then only do so for a short time.

If there is no lymphoedema in the 'at risk' arm, it is safe to put in a cannula or use a blood pressure cuff on the affected arm.



If you have had glands removed from both sides of your body, your anaesthetist will usually put a cannula in your non-dominant arm (the one you don't usually write with) or sometimes may put one in your foot.

You should discuss these risks and the best options for you with your anaesthetist.

Further information on lymphoedema can be found via the links below:

Macmillan Cancer Support (<u>bit.ly/3bfqlEU</u>) Lymphoedema Support Network (<u>lymphoedema.org</u>)



## Tell us what you think

We welcome suggestions to improve this leaflet. Please complete this short survey at: surveymonkey.co.uk/r/factsheetstest Or by scanning this QR code with your mobile:



### Disclaimer

We try very hard to keep the information in this leaflet accurate and up-to-date, but we cannot guarantee this. We don't expect this general information to cover all the questions you might have or to deal with everything that might be important to you. You should discuss your choices and any worries you have with your medical team, using this leaflet as a guide. This leaflet on its own should not be treated as advice. It cannot be used for any commercial or business purpose.

For full details, please see our website: rcoa.ac.uk/patientinfo/resources#disclaimer

#### Information for healthcare professionals on printing this leaflet

Please consider the visual impairments of patients when printing or photocopying this leaflet. Photocopies of photocopies are discouraged as these tend to be low quality prints and can be very difficult for patients to read. Please also make sure that you use the latest version of this leaflet, which is available on the RCoA website: <u>rcoa.ac.uk/patientinfo/factsheets</u>

#### **Royal College of Anaesthetists**

Churchill House, 35 Red Lion Square, London WC1R 4SG 020 7092 1500

### rcoa.ac.uk



#### Second Edition, February 2021

This leaflet will be reviewed within three years of the date of publication.

© 2021 Royal College of Anaesthetists

This leaflet may be copied for the purpose of producing patient information materials. Please quote this original source. If you wish to use part of this leaflet in another publication, suitable acknowledgement must be given and the logos, branding, images and icons removed. For more information, please contact us.

If you have any general comments, please email them to: patientinformation@rcoa.ac.uk