

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services (GPAS)

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

Consultation Draft November 2022



NICE has accredited the process used by the Royal College of Anaesthetists to produce its Guidance on the Provision of Anaesthesia Services. Accreditation is valid for five years from 2023.

More information on accreditation can be viewed at [www.nice.org.uk/accreditation](http://www.nice.org.uk/accreditation).

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 1 Authors

Dr Andrew Lindley  
Consultant Anaesthetist  
Leeds Teaching Hospital NHS Trust

Dr Prashant Kakodkar  
Consultant Anaesthetist  
Northampton General Hospital NHS Trust

Dr Iosifina Karmaniolou  
Locum Consultant in Anaesthetics  
Homerton University Hospital NHS Foundation Trust

Dr Baskar Manickam  
Consultant Anaesthetist  
Darlington Memorial Hospital  
County Durham and Darlington NHS  
Foundation Trust

Dr Subash Sivasubramaniam  
Consultant Anaesthetist  
Sandwell and West Birmingham Hospitals NHS Trust

2  
3  
4

### Chapter Development Group members

Mr Mike Donnellon  
Chair of the Education and Standards Committee  
The College of Operating Department Practitioners

Mr Robert Dudgeon  
Lay Representative, Lay Committee  
Royal College of Anaesthetists

Dr Nathaniel Duke  
Anaesthetist in Training  
University Hospitals Sussex NHS Foundation Trust

Mr William Eardley  
BOA Trauma Committee Vice-Chair  
South Tees Hospital  
British Orthopaedic Association

Dr Daniel Horner  
Consultant in Emergency Medicine and Intensive  
Care  
Royal College of Emergency Medicine

Dr Sunil Kumar  
SAS Lead, RCoA Council  
Royal College of Anaesthetists

Dr Purnima Narasimhan  
Anaesthetist in Training  
Queen Elizabeth Hospital Birmingham

5  
6  
7  
8  
9

### Acknowledgements

#### Peer reviewers

Dr Rajneesh Sachdeva  
Consultant Anaesthetist  
University Hospitals Birmingham NHS Foundation  
Trust

Dr Marcus Fletcher  
Consultant Anaesthetist  
Oxford Radcliffe Hospitals NHS Trust

10  
11  
12  
13  
14  
15  
16

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

17  
18  
19

### Chapter development technical team

Dr Rachel Evley  
Senior Research Fellow  
University of Nottingham

Ms Stephanie James  
Royal College of Anaesthetists

Dr Jeremy Langton  
GPAS Editor

Ms Ruth Nichols  
Royal College of Anaesthetists

Ms Zviko Nuamah  
Royal College of Anaesthetists

20

### Declarations of interest

21  
22  
23  
24  
25  
26  
27

All chapter development group (CDG) members, stakeholders and external peer reviewers were asked to declare any pecuniary or non-pecuniary conflict of interest, in line with the guidelines for the provision of anaesthetic services (GPAS) conflict of interest policy as described in the GPAS chapter development process document.

28  
29  
30  
31  
32

The nature of the involvement in all declarations made was not determined as being a risk to the transparency or impartiality of the chapter development. Where a member was conflicted in relation to a particular piece of evidence, they were asked to declare this and then, if necessary, remove themselves from the discussion of that particular piece of evidence and any recommendation pertaining to it.

### Medicolegal implications of GPAS guidelines

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

*GPAS guidelines are not intended to be construed or to serve as a standard of clinical care. Standards of care are determined based on all clinical data available for an individual case and are subject to change as scientific knowledge and technology advance and patterns of care evolve. Adherence to guideline recommendations will not ensure successful outcome in every case, nor should they be construed as including all proper methods of care or excluding other acceptable methods of care aimed at the same results. The ultimate judgement must be made by the appropriate healthcare professional(s) responsible for clinical decisions regarding a particular clinical procedure or treatment plan. This judgement should only be arrived at following discussion of the options with the patient, covering the diagnostic and treatment choices available. It is advised, however, that significant departures from the national guideline or any local guidelines derived from it should be fully documented in the patient's case notes at the time the relevant decision is taken.*

### Promoting equality and addressing health inequalities

46  
47  
48

The Royal College of Anaesthetists (RCoA) is committed to promoting equality and addressing health inequalities. Throughout the development of these guidelines we have:

49  
50  
51  
52  
53  
54  
55

- given due regard to the need to eliminate discrimination, harassment and victimisation, to advance equality of opportunity, and to foster good relations between people who share a relevant Protected Characteristic (as defined in the Equality Act 2010) and those who do not share it
- given regard to the need to reduce inequalities between patients in access to, and outcomes from healthcare services and to ensure services are provided in an integrated way where this might reduce health inequalities.

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 56 **GPAS Guidelines in context**

57 The GPAS documents should be viewed as 'living documents'. The GPAS guidelines development,  
58 implementation and review should be seen not as a linear process, but as a cycle of  
59 interdependent activities. These in turn are part of a range of activities to translate evidence into  
60 practice, set standards and promote clinical excellence in patient care.

61 Each of the GPAS chapters should be seen as independent but interlinked documents. Guidelines  
62 on the general provision of anaesthetic services are detailed in the following chapters:

- 63 • [Chapter 1: Guidelines for the Provision of Anaesthesia Services: The Good department](#)
- 64 • [Chapter 2: Guidelines for the Provision of Anaesthesia Services for the Perioperative Care of](#)  
65 [Elective and Urgent Care Patients.](#)

66 These guidelines apply to all patients who require anaesthesia or sedation, and are under the care  
67 of an anaesthetist. For urgent or immediate emergency interventions, this guidance may need to  
68 be modified as described in [Chapter 5: Guidelines for the Provision of Emergency Anaesthesia.](#)

69 The rest of the chapters of GPAS apply only to the population groups and settings outlined in the  
70 'Scope' section of these chapters. They outline guidance that is additional, different or particularly  
71 important to those population groups and settings included in the 'Scope'. Unless otherwise stated  
72 within the chapter, the recommendations outlined in chapters 1–5 still apply.

73 Each chapter will undergo yearly review, and will be continuously updated in the light of new  
74 evidence.

75 Guidelines alone will not result in better treatment and care for patients. Local and national  
76 implementation is crucial for changes in practice necessary for improvements in treatment and  
77 patient care.

### 78 **Aims and objectives**

79 The objective of this chapter is to promote current best practice for service provision in anaesthesia  
80 services for trauma and orthopaedic surgery. The guidance is intended for use by anaesthetists with  
81 responsibilities for trauma and orthopaedic surgery and healthcare managers.

82 This guideline does not comprehensively describe clinical best practice in anaesthesia for trauma  
83 and orthopaedic surgery, but is primarily concerned with the requirements for the provision of a  
84 safe, effective, well-led service. This service may be delivered by many different acceptable  
85 models. The guidance on provision of anaesthesia services for trauma and orthopaedic surgery  
86 applies to all settings where this is undertaken, regardless of funding. All age groups are included  
87 within the guidance unless otherwise stated, reflecting the broad nature of this service.

88 A wide range of evidence has been rigorously reviewed during the production of this chapter,  
89 including recommendations from peer-reviewed publications and national guidance where  
90 available. However, both the authors and the CDG agreed that there is a paucity of level 1  
91 evidence relating to service provision in anaesthesia services for trauma and orthopaedic surgery.  
92 In some cases, it has been necessary to include recommendations of good practice based on the  
93 clinical experience of the CDG. We hope that this document will act as a stimulus to future  
94 research.

95 The recommendations in this chapter will support the RCoA's Anaesthesia Clinical Services  
96 Accreditation (ACSA) process.

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 97 **Scope**

### 98 **Target audience**

99 All staff groups working in trauma, those attending trauma calls and orthopaedic surgery, including  
100 (but not restricted to) anaesthetists, operating department practitioners (ODPs), anaesthesia  
101 associates (AAs), nurses, allied health professionals and pharmacy staff.

### 102 **Target population**

103 All ages of patients undergoing trauma and orthopaedic surgery.

### 104 **Healthcare setting**

105 All settings within the hospital in which anaesthesia services for trauma and orthopaedic surgery are  
106 provided.

### 107 **Clinical management**

108 Key components needed to ensure provision of high-quality anaesthetic services for trauma and  
109 orthopaedic surgery.

110 Areas of provision considered:

- 111 • organisation, staffing and administration including leadership, clinical governance,  
112 policies, major trauma, emergency orthopaedics and elective orthopaedics
- 113 • areas of special requirements including children, pregnant patients and older patients
- 114 • equipment, services and facilities
- 115 • training and education
- 116 • research, audit and quality improvement
- 117 • patient information.

### 118 **Exclusions**

- 119 • Provision of trauma and orthopaedic surgery services provided by a specialty other than  
120 anaesthesia
- 121 • clinical guidelines specifying how healthcare professionals should care for patients
- 122 • national level issues.

123 General provision of critical care is outside the scope of this document. Further information,  
124 including definitions of levels of critical care can be found in the Faculty of Intensive Care Medicine  
125 and Intensive Care Society publication, [Guidelines for the Provision of Intensive Care Services 2022](#).

### 126 **Introduction**

127 Trauma remains the most common cause of loss of life in the under 40s age group in the UK, and as  
128 such major trauma centres (MTCs) and trauma units (TUs) have been established to receive  
129 patients of all ages, and improve outcomes.<sup>1</sup> Early anaesthetic involvement is beneficial at all  
130 stages, from the prehospital setting, to emergency departments (ED), operating rooms,  
131 interventional radiology suites, postoperative care units and the critical care environment. The  
132 need for significant anaesthetic input and support for these complex patients is an integral part of  
133 this pathway. Definitive fixation of all indicated fractures should be completed at the earliest  
134 possible opportunity.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

135 Hip fracture is the most common condition presenting for emergency orthopaedic surgery in the UK  
136 with many patients aged over 65. These patients along with those requiring surgical intervention for  
137 fragility fractures present significant challenges and the input from a multidisciplinary team and  
138 early surgery is essential to achieve good outcomes in this population.

139 Primary arthroplasty surgery significantly improves the quality of life and the mobility of those  
140 affected. With the advancing age of our population and their increasing expectations, the number  
141 of patients requiring primary arthroplasty surgery and subsequent revision arthroplasty surgery  
142 continues to rise. This population is frequently older with co-existing medical conditions that need to  
143 be optimised prior to surgery, and benefits from a multidisciplinary team (MDT) approach and the  
144 use of standardised protocols.

145 Orthopaedic surgery in children is wide ranging. From closed fracture manipulation and casting, to  
146 complex long bone and soft tissue surgery or spine correction of congenital or acquired conditions.  
147 These may be associated with neurological conditions, or specific syndromes that could pose  
148 challenges to those providing anaesthesia care.

### 149 **Recommendations**

150 The grade of evidence and the overall strength of each recommendation are tabulated in  
151 Appendix 1.

### 152 **1 Organisation, staffing and administration**

#### 153 **Medical leadership**

154 **1.1** Every department of anaesthesia should have a designated clinical lead (see [Glossary](#)) for  
155 anaesthesia services for trauma and a designated lead for anaesthesia services for  
156 orthopaedic surgery. This should be recognised in their job plan and they should be involved  
157 in multidisciplinary service planning and governance within the unit. They should have access  
158 to trust board through a governance structure with explicit pathways of communication.

159 **1.2** Autonomously practising anaesthetists and intensivists should be involved in the planning of  
160 local trauma services. Those with defined responsibility for major trauma management should  
161 be engaged in the layout and logistics of the resuscitation room, interventional radiology and  
162 theatres suites.

163 **1.3** Each department of anaesthesia should have an annual plan in place for the workload to be  
164 delivered safely and effectively.<sup>2</sup>

165 **1.4** Organisations should explicitly recognise the 24/7 nature of trauma work, and this requires a  
166 specific organisational approach for standards to be achieved throughout the whole of the  
167 week.

168 **1.5** The provision of a high-quality trauma and orthopaedic service should be an explicit aim of  
169 the hospital executive and senior staff team. This should be reflected in hospital published  
170 plans and by the provision of a management structure to support this aim.<sup>3,4,5</sup>

171 **1.6** Hospital business plans should address the predicted growth in elective and non-elective  
172 demands due to expanding local population and an ageing population.<sup>6</sup> Planning should be  
173 based on accurate and timely demand and capacity modelling.<sup>7</sup>

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 174 **Clinical Governance**

175 Clinical Governance is covered in detail in [Chapter 1: Guidelines for the provision of Anaesthesia](#)  
176 [Services: The Good Department](#). The principles of governance described in the chapter are  
177 applicable to provision of services for trauma and orthopaedic surgery.

178 **1.7** Regular multidisciplinary mortality and morbidity meetings should take place in all trauma  
179 centres and follow the guidance of the World Health Organization (WHO).<sup>8</sup>

180 **1.8** Anaesthetists should be involved in multidisciplinary governance meetings. Perioperative  
181 outcome data should be discussed in these meetings.

182 **1.9** Care pathways and care bundles for common procedures such as hip fracture improve  
183 outcomes. Anaesthetists should be involved in developing, delivering and evolving these  
184 pathways and bundles.

185 **1.10** Governance meetings should take place across the entire trauma network at defined  
186 intervals. Besides individual case discussion, feedback information from the Trauma Audit and  
187 Research Network (TARN) should be disseminated, and mechanisms set in place to correct  
188 any problems identified.<sup>1</sup>

### 189 **Policies**

190 General policies detailed in [Chapter 2: Guidelines for the provision of Anaesthesia Services for](#)  
191 [Perioperative Care of Elective and Urgent Care patients](#) are relevant to provision of anaesthesia  
192 services for trauma and orthopaedic surgery.

193 **1.11** Specific local policies pertinent to trauma and orthopaedic surgery should be developed by  
194 a multidisciplinary team including an anaesthetist, acute pain nurse, pharmacist,  
195 physiotherapist, critical care clinicians, surgeons and other relevant specialties.<sup>9</sup>

196 **1.12** Local policies should be in agreement with relevant published national guidelines.

197 **1.13** Local policies should be easily accessible to all staff caring for trauma and orthopaedic  
198 patients. These include but are not limited to:

- 199 • preoperative screening for complex pain issues and access to acute pain services and  
200 advanced pain management methods
- 201 • pain management pathways for patients with chest injuries
- 202 • supervision and monitoring of patients by competent clinical staff during surgery  
203 performed under peripheral nerve blocks, including in a block room or similar facility
- 204 • perioperative anticoagulation guidelines for safe placement of epidural and regional  
205 nerve block techniques<sup>10</sup>
- 206 • recognition and management of patients at risk of acute compartment syndrome<sup>11</sup>
- 207 • 'Stop Before You Block'<sup>12,13,14,15</sup>
- 208 • management of complications of regional anaesthesia including high spinal block and  
209 accidental dural puncture
- 210 • assessment and management of local anaesthetic systemic toxicity
- 211 • assessment and management of peripheral nerve injury
- 212 • post procedure monitoring of epidurals, nerve blocks and continuous infusion analgesia on  
213 the ward, including follow up care in hospital and after discharge
- 214 • protocol for whom to call for problems with postoperative pain relief

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 215 • venous thromboembolism prophylaxis after orthopaedic and trauma surgery.<sup>16</sup>
- 216 1.14 Units should use a delirium assessment tool and have a delirium policy describing  
217 preventative measures and its management.<sup>17</sup>
- 218 1.15 There should be specific guidelines for assessing a suspected difficult airway in patients with  
219 spine and joint disease and for measuring lung function in patients with kyphoscoliosis.<sup>18</sup>
- 220 1.16 Patients at risk of acute compartment syndrome (ACS) should be identified on admission to  
221 hospital or at the time of surgery, and should be managed within agreed, multidisciplinary  
222 protocols.<sup>19</sup>
- 223 **2 Major trauma**
- 224 **Policies**
- 225 2.1 Local policies should ensure National Institute for Health and Care Excellence (NICE)  
226 recommendations and quality standards for major trauma services are met and should be  
227 agreed with the regional trauma network.<sup>20,21</sup>
- 228 2.2 There should be local policies in place to meet the quality standards developed by NICE for:<sup>21</sup>
- 229 • patients with major trauma who cannot maintain their airway and/or ventilation to have  
230 drug-assisted rapid sequence induction (RSI) of anaesthesia and intubation within  
231 45 minutes of the initial call to the emergency services
- 232 • patients who have had urgent 3D imaging for major trauma have a provisional written  
233 radiology report within 60 minutes of the scan
- 234 • patients with open fractures of long bones, the hindfoot or midfoot have fixation and  
235 definitive soft tissue cover within 72 hours of injury if this cannot be performed at the same  
236 time as debridement
- 237 • patients with full in-line spinal immobilisation to have their risk of cervical spine injury  
238 assessed using the Canadian C-spine rule<sup>22</sup>
- 239 • major trauma centres to have a dedicated trauma ward for patients with multisystem  
240 injuries and a designated consultant available to contact 24 hours a day, 7 days a week.
- 241 • major trauma centres to have acute specialist services for rehabilitation after major  
242 trauma, and for children and older people.
- 243 2.3 Initial management should follow the adult trauma life support principles with management  
244 of airway, breathing and circulation along with cervical spine stabilisation occurring in  
245 parallel rather than in sequence.<sup>23</sup> Local guidelines should be followed to ensure the  
246 appropriate tier of trauma call response is made.
- 247 2.4 Pain management pathways should be followed for chest wall injuries including provision for  
248 early epidural or nerve blocks in patients with multiple rib fractures.<sup>24</sup>
- 249 2.5 Assessment and management for a cervical spine injury should follow pre agreed existing  
250 NICE and British Orthopaedic Association guidance.<sup>25,26</sup> Spinal clearance protocols should be  
251 embedded into practice.<sup>27</sup>
- 252 2.6 There should be a local protocol in place for emergency access to an operating theatre or  
253 intervention suite, to provide rapid intervention in life threatening or limb threatening  
254 conditions.<sup>28</sup>
- 255 2.7 All acute hospitals should have a defined major incident plan. The plan should be built  
256 around the regional network of MTCs, TUs and local emergency hospitals (LEHs).<sup>20</sup>

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 257 **Organisation**

258 2.8 A structured system for recording and receiving information about trauma patients should be  
259 developed and implemented. Clear lines of communication to establish this prehospital  
260 documentation should be shared with senior nursing staff and trauma teams.<sup>20,29</sup>

261 2.9 Rapid and effective communication is crucial in emergency situations. Communication  
262 strategies should consider the use of technologies such as smart phones, and standardised  
263 methodology such as situation, background, assessment, recommendation.<sup>30</sup>

264 2.10 Patients who have sustained rib fractures should have an early multidisciplinary assessment  
265 involving multidisciplinary teams such as surgeons, pain services, critical care and  
266 physiotherapy to determine the optimal analgesia and definitive management options to  
267 minimise complications related to altered pulmonary mechanics, lung capacity and  
268 ventilation.

269 2.11 There should be a flexible approach to trauma list planning and management to  
270 accommodate emergency cases which need priority treatment. There should be a system in  
271 place to alert the theatre team of the arrival of an unstable patient with major trauma.  
272 Appropriately trained staff and facilities should be available to receive these patients at short  
273 notice.<sup>18</sup>

274 2.12 The trauma team should attend to all cases of suspected major trauma, according to the  
275 predefined local criteria. The trauma team should also be present for paediatric and older  
276 patients (where appropriate), patients with unexpected findings on arrival and to receive  
277 patients following interhospital transfer.

278 2.13 The time interval between the injury and initiation of management should be considered for  
279 patients who have sustained trauma to improve patient outcomes.

280 2.14 The optimal destination for the majority of patients with major trauma is a major trauma  
281 centre (MTC). A pre-hospital triage tool should be used to differentiate between patients who  
282 should be diverted to a MTC and those who could be taken initially to a trauma unit (TU) for  
283 treatment.<sup>20,31</sup> Major trauma patients should only be taken to a TU if the patient needs a life-  
284 saving intervention that cannot be delivered by the pre-hospital team.

285 2.15 MTCs should have a clear point of contact to provide clinical advice via clearly identified  
286 pathways and local tier arrangement to other providers within the network. This includes  
287 advice during the pre-hospital stage and whilst patients are awaiting transfer to MTCs for  
288 definitive treatment.

289 2.16 The anaesthetist plays a key role in the multidisciplinary team who apart from being involved  
290 in airway management in major trauma patients, should provide input into the recognition  
291 and management of acute physiological derangement, haemorrhage, and shock.

292 2.17 Handovers for patients requiring emergency trauma surgery should be structured to ensure  
293 continuity of care. Handover protocols should include clear documentation of care delivered  
294 and the future treatment plan for the patient.<sup>18,32</sup>

### 295 **Staffing**

296 2.18 Anaesthesia for the emergency control of major traumatic haemorrhage, and other damage  
297 limiting interventions in the operating theatre or radiology intervention suite, should be  
298 consultant anaesthetist led. Where consultants are not resident, clear lines of communication  
299 and notification should be in place to allow early attendance to trauma calls.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

300 2.19 A major trauma coordinator (or someone in a similar role) should be responsible for  
301 overseeing scheduled admissions, preparation and planning for surgery, and coordination of  
302 care of trauma patients with other specialties including critical care and postoperative care.

### 303 **Equipment, services and facilities**

304 2.20 Immobilisation equipment including a range of appropriately sized semi rigid collars, head  
305 blocks, tape, a vacuum mattress and a scoop board should be available. Spinal clearance  
306 should be achieved as soon as clinically possible, to minimise discomfort and complications  
307 from prolonged immobilisation in patients who do not have spinal injuries.

308 2.21 In suspected spinal injury, hard spinal boards should only be used as a prehospital extrication  
309 device and not be used for transport.<sup>33</sup> A scoop stretcher or full-length vacuum mattress  
310 should be used for transfer.

311 2.22 Equipment for portable monitoring to level 3 standards and ventilation should be available in  
312 the ED resuscitation room.<sup>34,35</sup>

313 2.23 Use of point of care ultrasound (POCUS) should be available as it is a useful adjunct to the  
314 primary survey in acute trauma. Operators need to ensure this does not delay time to  
315 definitive imaging or intervention.<sup>36,37,38</sup>

316 2.24 Patients who have acute nerve or spinal cord compression should be referred immediately to  
317 a neurosurgeon or specialist spinal unit (where required).<sup>39</sup>

318 2.25 In MTCs with a high volume of patients, prethawed plasma should be immediately available.

319 2.26 For patients with complex trauma, including spinal cord injuries and traumatic brain injury,  
320 there should be rapid access to key professionals and regional specialists. Patients, relatives  
321 and carers should be directed to appropriate support groups where relevant e.g. the Spinal  
322 Injuries Association.<sup>40</sup>

323 2.27 An emergency operating theatre should be rapidly available at all times for major trauma  
324 patients. The available equipment should be suitable for a full range of emergency trauma  
325 procedures. Use of this theatre for non-urgent cases should be tightly controlled. If the  
326 designated emergency theatre is occupied, there should be a robust, flexible and agreed  
327 backup plan to obtain an appropriate alternative theatre for the next emergency case.

328 2.28 The emergency operating theatre should be equipped with a radiolucent operating table  
329 that allows fluoroscopic imaging of all body parts without repositioning the patient.

330 2.29 In MTCs and TUs there should be a rapidly accessible imaging suite for patients with major  
331 trauma, which has immediate access to specialised equipment for the management of  
332 difficult airways including physiological and gas monitoring. In addition, the room design  
333 should allow visual and technical monitoring of the patient by the anaesthetic staff.<sup>41</sup>

334 2.30 In MTCs and TUs, the resuscitation room receiving bays should be large enough to allow  
335 simultaneous emergency procedures to be performed by trauma team members.

336 2.31 Hospitals admitting patients with major trauma should have critical care to both Level 2 and 3  
337 standards on site.<sup>42</sup> Portable invasive haemodynamic monitoring should be available to  
338 facilitate transfer to and from the critical care areas.

### 339 **Transportation of the trauma patient**

340 2.32 When transporting a trauma patient, the following should be available:

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 341
  - appropriately trained and competent staff
- 342
  - insurance (personal and medical indemnity)
- 343
  - crash test compliant equipment
- 344
  - ambulance booking procedures
- 345
  - procedures for receiving patients
- 346
  - communication between medical teams and families
- 347
  - documentation and procedures for repatriation of staff and equipment once the transfer
- 348
  - and handover are completed.<sup>9,43,44</sup>
- 349 **2.33** Transport of patients within the hospital and between hospitals (e.g. transfers to major trauma,
- 350 neurosurgical or paediatric centres) should be undertaken in a timely manner, without
- 351 unnecessary delays, and in accordance with established guidelines and standards.<sup>9,43,44,45,46</sup>
- 352 Hospital transfers may involve a retrieval service.
- 353 **2.34** Local guidelines for patient transfers between referring hospitals, neurosurgical units and local
- 354 ambulance services should be consistent with national guidelines for the safe transfer of the
- 355 brain-injured patient.<sup>47</sup>
- 356 **3 Emergency orthopaedics**
- 357 **Fragility hip fracture**
- 358 **3.1** For patients who have sustained hip fractures, femoral or fascia iliaca nerve blocks should be
- 359 provided in the emergency department (ED) and at the time of surgery (provided six hours
- 360 has passed between blocks).<sup>48</sup>
- 361 **3.2** Hospitals providing surgical treatment for hip fractures should have a formal pathway
- 362 including prompt provision of analgesia (including nerve blocks) and hydration, preoperative
- 363 assessment of high-risk patients by the anaesthetic team, along with, orthogeriatrician input
- 364 and be prioritised on orthopaedic trauma lists.<sup>49,50,51,52</sup>
- 365 **3.3** Risk assessment should be performed in all patients with hip fracture. The Nottingham Hip
- 366 Fracture score and National Hip Fracture Database Tool could be used to assess risk.<sup>53</sup> Frailty
- 367 scores, 4 'A's test score for delirium, Nottingham Hip Fracture Risk Score for Kidney Injury are
- 368 useful organ specific assessment tools.
- 369 **3.4** Anaesthetists should facilitate surgery within 36 hours of a hip fracture.<sup>54</sup> Surgery should be
- 370 delayed only if the benefits of additional medical treatment outweigh the risks of delaying
- 371 surgery. The risks of delay associated with pain and immobility contribute to poor outcomes to
- 372 a far greater extent than correction of an abnormality to a particular numerical value.<sup>48</sup>
- 373 **3.5** Dedicated trauma operating lists should be scheduled daily, including weekends to meet
- 374 local demands, and ensure 36 hours targets for hip fracture are met. Extra provision during
- 375 the day and in the evenings may be necessary to meet local demands and limit overnight
- 376 operating.<sup>55</sup>
- 377 **3.6** Unoperated hip fractures in older patients have a high mortality rate. Evidence shows ASA4
- 378 patients have a higher survival rate when managed surgically.<sup>56</sup> Hip fracture surgery should
- 379 be considered for patients even in the presence of significant comorbidities. Provision for safe
- 380 anaesthesia and recovery of these patients, including handover to ward teams, should be
- 381 available to facilitate this.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 382 3.7 Perioperative anaesthetic care for a patient who is older with a hip fracture should be  
383 standardised with the overarching goals of management being patient remobilisation, re-  
384 enablement and rehabilitation
- 385 3.8 Anaesthesia (and surgery) for hip fractures should be undertaken by an appropriately  
386 experienced anaesthetist (and surgeon).
- 387 3.9 An anaesthetic job planned representative should attend regular multidisciplinary hip  
388 fracture management meetings and feedback any relevant learning points to departments  
389 of anaesthesia and individual anaesthetists, as appropriate.
- 390 3.10 Facilities to provide total hip replacement to hip fracture patients with limited comorbidities  
391 should be available seven days a week.
- 392 3.11 Safe protocols procedure kits and antidotes should be readily available, to adopt best  
393 practice and allow timely delivery of fascia iliaca compartment blocks for hip fracture in the  
394 ED.<sup>57</sup>
- 395 3.12 There should be a formalised integrated pathway for high-risk trauma patients, such as hip  
396 fractures, which includes:<sup>4,33,58,59</sup>
- 397 • a clear diagnostic and management plan made on admission<sup>56</sup>
  - 398 • a clear identification and documentation of comorbidities
  - 399 • a clear preoperative assessment and optimization plan by an anaesthetist and/or an  
400 orthogeriatrician<sup>60</sup>
  - 401 • documentation of preoperative investigations and testing
  - 402 • a reconciled list of their medicines to assess the risk of existing medications (including  
403 anticoagulation) and the risk associated with stopping long term medication<sup>56,57,61</sup>
  - 404 • documentation of risk assessment, mortality risk, discussions with family and any other  
405 important decisions<sup>25,33,62</sup>
  - 406 • informed consent for surgery including identification of decision-making proxies i.e. a  
407 lasting power of attorney<sup>4,63</sup>
  - 408 • a plan for postoperative care.<sup>4,63</sup>
- 409 3.13 Agreed local guidelines should be in place and implemented on the following:
- 410 • compliance with best practice anaesthetic management protocols for hip fracture as  
411 recommended by the Association of Anaesthetists.<sup>49,64</sup>
  - 412 • tailored World Health Organization (WHO) safety checklists to discuss the requirement for  
413 use of bone cement
  - 414 • preoperative assessment for treatment escalation and cardiopulmonary resuscitation
  - 415 • older people (>65 years) and/or frail people with long bone and periprosthetic fractures  
416 should receive similar treatment as those with hip fracture.
- 417 3.14 Departments should develop protocols for reviewing hip fracture patients postoperatively, in  
418 order to support ongoing orthogeriatric care, and also to learn from successes and problems  
419 as part of continuous quality improvement.<sup>48</sup>

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 420 **4 Elective orthopaedics**

#### 421 **Organisation and administration**

422 Detailed recommendations for organisation and administration of anaesthesia services for elective  
423 surgery are detailed in [Chapter 2: Guidelines for the provision of Anaesthesia Services for](#)  
424 [Perioperative Care of Elective and Urgent Care patients](#).

425 4.1 There should be a preoperative assessment clinic for elective orthopaedic surgery.

426 4.2 The anaesthetist should contribute in the multidisciplinary perioperative care process which  
427 focuses on preoptimisation, patient education, standardised enhanced recovery pathways  
428 of care aimed at delivering early mobility, discharge, and early return to normal life.<sup>65,66</sup> The  
429 option of doing nothing should be considered where relevant.<sup>67,68</sup>

430 4.3 There should be multidisciplinary input for the preoperative assessment of high risk patients  
431 such as patients with cognitive disorders, chronic kidney disease, diabetes mellitus and  
432 ischaemic heart disease.<sup>69,50</sup> The anaesthetist should be involved in preoperative optimisation  
433 and prehabilitation plans.<sup>66,63</sup>

434 4.4 In patients aged over 65, frailty screening using an appropriate validated screening tool  
435 should be performed and documented early in the preassessment pathway. A screening tool  
436 used in combination with direct questioning should also be adopted to help identify patients  
437 with cognitive impairment and therefore increased risk of delirium.<sup>70</sup>

438 4.5 Patients should be screened for chronic pain and opioid use in the preoperative period.  
439 Preoptimisation should ensure optimal management of preoperative pain, psychological  
440 preparation, education, and expectation management.<sup>52,71</sup>

441 4.6 A perioperative management plan should be formulated for all patients and should include  
442 multimodal analgesia and intrathecal opioid sparing analgesic techniques.<sup>72</sup> Multimodal  
443 analgesic techniques should aim to provide optimal pain relief whilst minimizing side effects  
444 such as sedation, postoperative nausea, and vomiting (PONV), and hypotension which might  
445 compromise early rehabilitation and recovery.<sup>71,73,74</sup>

446 4.7 There should be an enhanced recovery after surgery (ERAS) programme for suitable patients  
447 undergoing elective orthopaedic surgery as it improves early mobilisation, reduces length of  
448 stay, postoperative complications and mortality.<sup>75</sup>

449 4.8 Elective patients with major comorbidities or those undergoing complex or prolonged surgery  
450 should be scheduled earlier in the day, to allow time for postoperative stabilisation.

451 4.9 Elective orthopaedic operating lists should be separated from trauma lists, to allow efficiency,  
452 ensure safety, prevent cancellations and enable a flexible response for emergencies.

453 4.10 Hospitals should consider providing specific regional anaesthesia lists and using dedicated  
454 areas for performing peripheral nerve blocks.<sup>76</sup>

455 4.11 Elective orthopaedic units performing major inpatient surgery should have 24/7 access to all  
456 support services including acute pain services and critical care.

### 457 **5 Equipment, facilities and services**

458 Detailed recommendations for equipment, facilities and services of anaesthesia services for  
459 elective surgery are detailed in [Chapter 2: Guidelines for the provision of Anaesthesia Services for](#)  
460 [Perioperative Care of Elective and Urgent Care patients](#).

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 461 **Equipment**

- 462 5.1 Elective orthopaedic and planned trauma cases should have their temperature checked  
463 preoperatively on the ward. Warming devices for patients should be available for use in the  
464 anaesthetic room, operating theatre, recovery unit and ED.<sup>3</sup>
- 465 5.2 A range of operating tables with attachments for spinal, thoracic, pelvic and limb trauma  
466 procedures should be available as appropriate.<sup>77</sup>
- 467 5.3 Tourniquets and inflation devices of suitable sizes should be available for upper and lower  
468 limb surgery requiring a bloodless field.
- 469 5.4 Cell salvage equipment should be available for cases where significant blood loss is  
470 anticipated.<sup>78,79</sup> Staff who operate this equipment should receive training in how to operate  
471 it, and use it with sufficient frequency to maintain their skills.
- 472 5.5 A rapid infuser allowing the infusion of warmed intravenous fluids and blood products should  
473 be available in the anaesthetic room, operating theatre, recovery unit and ED.<sup>80</sup>
- 474 5.6 Equipment to facilitate haemodynamic and cardiac output monitoring should be available.
- 475 5.7 A standardised 'difficult airway trolley' should be immediately available in all areas where  
476 major trauma patients are received or managed to level 2 (or greater) standards of care.  
477 These should be equipped as defined in the Difficult Airway Society (DAS) guidelines.<sup>5,81,82</sup>
- 478 5.8 Patient positioning for elective and trauma orthopaedic surgery involves a variety of specialist  
479 equipment, tables and attachments. These should be suitable to manage patients across a  
480 wide weight range, with theatre personnel aware of the upper weight limits.<sup>83,84</sup>

### 481 **Facilities**

- 482 5.9 Primary and revision arthroplasty surgery, along with trauma surgery involving bone implants  
483 or internal fixation should be carried out in an operating theatre with multiple air changes per  
484 hour (e.g., laminar flow).
- 485 5.10 Where appropriate, point of care testing for haemoglobin, blood gases, lactate, ketones,  
486 coagulation and blood sugar should be available for patients with major trauma throughout  
487 the patient journey and those undergoing orthopaedic procedures associated with a risk of  
488 haemorrhage.<sup>85</sup> If near-patient testing is not available, laboratory testing should be readily  
489 and promptly available.<sup>86</sup>
- 490 5.11 Transport and distribution of blood and blood components at all stages of the transfusion  
491 chain must be maintained under appropriate conditions to ensure the integrity of the  
492 product.<sup>87</sup>
- 493 5.12 Appropriate blood storage facilities should be clearly identified and provided in close  
494 proximity to the emergency operating theatre.<sup>88</sup>
- 495 5.13 Tranexamic acid should be available for administration if major haemorrhage is suspected in  
496 a trauma patient within three hours of injury.<sup>89,90,91</sup>
- 497 5.14 Newer hemostatic agents and antidotes should be readily available to clinical teams for  
498 emergency and trauma surgery.<sup>92</sup>
- 499 5.15 There should be adequate provision of postoperative beds for orthopaedic and trauma  
500 patients to allow timely discharge of patients from theatre recovery areas.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

501 5.16 A fully equipped high dependency unit (HDU) of Level 2 standards should be available on site  
502 for high-risk patients undergoing major orthopaedic surgery, including revision joint  
503 replacement and surgery involving instrumentation of the spine. If the hospital does not have  
504 a Level 3 facility, protocols should be in place to determine when and how to transfer to a  
505 hospital with a Level 3 facility. Alternatively, other models of enhanced postoperative care  
506 such as a post anaesthesia care unit (PACU) should be considered.

507 5.17 Clinical deterioration can be identified using early warning scores and mitigated by  
508 proactively reviewing patients at risk. Arrangements to seek critical care input in deteriorating  
509 patients in a timely manner should be in place.

## 510 6 Areas of Special Requirements

### 511 Children

512 Anaesthetists will often be part of the MDT responsible for the initial resuscitation and stabilisation of  
513 the critically ill or injured child, prior to transfer to a specialist centre. All hospitals with an ED will be  
514 exposed to high volumes of paediatric patients with low energy transfer injuries. Detailed  
515 recommendations for paediatric services are described in [GPAS Chapter 10: Guidelines for the](#)  
516 [Provision of Paediatric Anaesthesia Services 2022](#).

517 6.1 Staff should be vigilant for non-accidental injury in children with trauma injuries and should  
518 make enquiries of the circumstances around major trauma and ask if there are safeguarding  
519 concerns.

520 6.2 Healthcare workers, including the anaesthetist, must be aware of the local policy for child  
521 protection, and that they have an obligation to document and report any concerns to a  
522 responsible individual.<sup>93</sup>

523 6.3 Hospitals must have guidelines in place to ensure the safety of children admitted to hospital,  
524 to monitor injured children known to be at risk, and identify concerns arising from any injury or  
525 pattern of injuries.<sup>94</sup> They must provide the appropriate training related to these guidelines.

### 526 Pregnant patients

527 Trauma is a leading cause of non-obstetric mortality in pregnant patients.<sup>4</sup> Although the primary  
528 duty of care is to the mother, fetal and maternal wellbeing are inextricably linked. Standards for  
529 non-obstetric emergency procedures in pregnant patients are described in [GPAS Chapter 5:](#)  
530 [Guidelines for the Provision of Emergency Anaesthesia Services](#).

531 6.4 A standardised approach should be taken when completing a trauma survey in pregnancy  
532 and a maternal-foetal trauma checklist should be considered as this lays the foundation for  
533 interdisciplinary collaboration in a stressful environment.<sup>95</sup>

534 6.5 Provision for fetal monitoring and emergency lower (uterine) segment caesarean section  
535 should be available if indicated in the ED.<sup>96,97,98</sup>

536 6.6 In cases of pregnant orthopaedic trauma patients, diversion to a centre with obstetric and  
537 trauma expertise directly from the scene of an injury should be considered, to avoid delay of  
538 appropriate specialist care.<sup>99</sup>

### 539 Older patients

540 6.7 A patient centred approach is preferred for documenting advanced care plans that include  
541 overall treatment goals including resuscitation status. It should include discussing and  
542 planning treatments that should be considered, not just those that should be withheld.<sup>18</sup>

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 543 6.8 A fall of <2m is the commonest mechanism of injury in older patients. Prehospital triage to aid  
544 early identification of severe injuries in older patients should be available to allow quick  
545 transfer from TU to a MTC for specialist investigation and intervention.<sup>100</sup>
- 546 6.9 Older patients who are admitted following trauma should have a comprehensive geriatric  
547 assessment. The use of frailty screening tools may facilitate more informed early decision  
548 making in older trauma patients.<sup>101</sup>
- 549 6.10 Protocols for end of life care should be in place to manage older patients with frailty who are  
550 unlikely to survive. The multidisciplinary team and patient's family or next of kin should be  
551 involved in these decisions.<sup>102</sup>
- 552 **7 Training and Education**
- 553 7.1 All patients undergoing anaesthesia should be under the care of an autonomously practising  
554 anaesthetist whose name is recorded as part of the anaesthetic record.<sup>4,103</sup>
- 555 7.2 All anaesthetists providing anaesthesia for trauma and orthopaedics should have  
556 appropriate knowledge, skills, attitudes and behaviour in accordance with the RCoA training  
557 standards.<sup>104</sup>
- 558 7.3 Anaesthetists with a specific interest in orthopaedics and trauma should deliver regular  
559 theatre sessions to ensure the maintenance of their skills and experience.
- 560 7.4 Anaesthetists with responsibility for the intraoperative care of trauma patients should ensure  
561 that their skills and knowledge of current recommendations are up to date, particularly in the  
562 management of major haemorrhage.
- 563 7.5 MTC and TU anaesthetic departments should consider appointing anaesthetists with an  
564 interest in prehospital care (see [Glossary](#)). Anaesthetists who provide prehospital care outside  
565 the hospital setting should be qualified to do so.<sup>105</sup>
- 566 7.6 Anaesthetists who manage patients with major trauma should consider undertaking  
567 advanced trauma life support (ATLS), European Trauma Course (ETC) or equivalent training,  
568 and should update their training at regular intervals.
- 569 7.7 Anaesthetists providing anaesthesia for trauma and orthopaedic surgery should learn and  
570 maintain expertise in a wide range of regional anaesthetic techniques, including central and  
571 peripheral neural blockade.<sup>104</sup>
- 572 7.8 All anaesthetists involved in the management of major trauma should understand the  
573 principles and techniques of haemorrhage control resuscitation to prevent the lethal  
574 diamond of hypothermia, acidosis, coagulopathy and hypocalcemia using low volume fluid  
575 resuscitation, blood products and damage control surgery.<sup>106,107,108</sup>
- 576 7.9 Appropriately trained theatre staff should be available when treating patients with multiple  
577 injuries. They should be skilled and experienced in all surgical specialties that may present in  
578 the treatment of patients with multiple injuries.
- 579 7.10 Anaesthetic trauma theatre teams should be trained in the correct use of all essential  
580 anaesthetic theatre equipment used for trauma surgery.
- 581 7.11 Staff in the recovery area and in the wards who receive patients after surgery with epidural  
582 infusions, nerve blocks or intravenous opioid infusions (including patient controlled analgesia)  
583 should have received up to date formal training in caring for such patients.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 584 7.12 Staff expected to top up medication for patients with epidurals and continuous nerve  
585 blockade should be trained in administration of such medications.
- 586 7.13 Anaesthetic practitioners involved in the administration of anticoagulant therapies should  
587 have current and up to date knowledge in their use.
- 588 7.14 There should be regular multidisciplinary in situ simulation training for the initial management  
589 of major trauma care and resuscitation to standardise clinical practice. Simulation can  
590 improve technical and non-technical skills including communication and teamwork.<sup>109</sup>
- 591 7.15 Appropriate early referral to the acute pain services and use of regional analgesia should be  
592 considered in patients with chest trauma.<sup>110</sup>
- 593 7.16 The diagnostic and therapeutic applications of POCUS in trauma are expanding. There  
594 should be emphasis on quality training and a robust governance process for POCUS  
595 operators within the trauma multidisciplinary team.<sup>111,112</sup>
- 596 7.17 Major incident training exercises should take place at regular intervals.
- 597 7.18 Organisations should provide mandatory training relating to advance care planning and  
598 resuscitation policies and documents.<sup>113</sup>
- 599 7.19 Educational opportunities for anaesthetists in training in MTC and TU will undoubtedly occur  
600 during predictable job planned consultant direct clinical care sessions out of hours due to the  
601 nature of trauma. Hospitals in which anaesthetists in training work a full or partial shift system  
602 should consider providing additional consultant programmed activities to allow training and  
603 supervision to take place in the evening.
- 604 7.20 Hospitals should consider training ED staff in acute pain management of both adult and  
605 paediatric patients with trauma, in particular using ultrasound guided femoral nerve block or  
606 fascia iliaca block for hip fractures in elderly patients and femoral fractures in children.<sup>114,115</sup>
- 607 7.21 The definitive care of complex spinal and pelvic injuries requires early multidisciplinary  
608 specialist spinal (orthopaedic or neurosurgical surgery) and pelvic team discussion. The  
609 anaesthetist managing such cases should have appropriate training and experience in  
610 management of these complex patients including management of associated  
611 complications.
- 612 **8 Research audit and quality improvement**
- 613 8.1 All sites should consider participating in active research studies on the National Institute for  
614 Health and Care Clinical Research Network portfolio for Trauma and Emergency Care.<sup>116</sup>
- 615 8.2 All major trauma centres should have a dedicated research lead with appropriate job  
616 planned time and should receive training on ethical and organisational issues.<sup>117</sup>
- 617 8.3 All clinicians involved in trauma care should be aware of active studies and consider  
618 completing good clinical practice training and participating in screening and recruiting of  
619 research participants.<sup>116,118</sup>
- 620 8.4 Opportunities for associate principal investigator roles should be encouraged.<sup>116</sup>
- 621 8.5 Trauma and orthopaedic surgery should be included in anaesthetic departmental audit  
622 programmes, including ongoing audit of complications and adverse events. The trauma  
623 anaesthetists should have provision in their job plan to attend trauma MDT meetings for  
624 discussion regarding high risk patients.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 625 8.6 All hospitals treating patients with hip fractures should participate in national audits, e.g.,  
626 National Hip Fracture Database or the National Joint Registry to monitor its performance  
627 against national benchmarks, quality standards, and contribute to research. Outcomes from  
628 these audits should be discussed at governance meetings and distributed to anaesthetic  
629 staff.<sup>119,120</sup>
- 630 8.7 All hospitals receiving major trauma cases should contribute to the Trauma Audit and  
631 Research Network (TARN), to monitor their performance against national benchmarks and  
632 quality standards and contribute to research.<sup>1</sup> Comparative data analysis and display on the  
633 national major trauma dashboard (via TARN) is invaluable for quality assurance.
- 634 8.8 MTCs and TUs in England should undergo regular peer reviews within the National Peer  
635 Review Programme with their performance judged according to national major trauma  
636 measures.<sup>121</sup>
- 637 8.9 All new spinal cord injury patients should be referred through the NHS Spinal Cord Injury  
638 Service (NSCIS) and registered on the National Spinal Cord Injury Database (NSCID).<sup>28</sup> The  
639 incidence of complications should be recorded.
- 640 8.10 There should be clear processes and policies for reporting and learning from near misses and  
641 critical incidents. National patient safety alerts should be communicated and actions agreed  
642 locally to reduce the risk of harm.
- 643 8.11 Nationally agreed key performance indicators should be used to monitor the performance of  
644 the pathways for hip fractures and major trauma and reviewed by a multidisciplinary  
645 committee including a Trauma Lead anaesthetist. In addition, local quality indicators should  
646 be developed proactively, to support continuing improvement of these services within  
647 organisations.
- 648 8.12 Impact of enhanced recovery pathways for elective surgery should be audited to focus  
649 beyond the length of stay to improve patient outcome and satisfaction.
- 650 8.13 Evaluation of patient centred outcomes on pain management and quality of recovery in  
651 hospital and after discharge using a validated questionnaire can be a useful tool to guide  
652 quality improvement in care pathways.<sup>122,123</sup>
- 653 8.14 Organisations should have a service improvement team that coordinates national and local  
654 projects, and encourages a multidisciplinary approach to trauma and orthopaedic services.  
655 Data should be collected to provide high quality information to drive change and support  
656 service development. Good data, Quality improvement tools and organisational support can  
657 create feedback strategies which drive improvement.<sup>124</sup>

## 658 9 Patient information

659 The Royal College of Anaesthetists have developed a range of [Trusted Information Creator](#)  
660 [Kitemark](#) accredited patient information resources that can be accessed from our [website](#). Our  
661 main leaflets are now translated into more than 20 languages, including Welsh.

### 662 Information for patients, relatives and carers

- 663 9.1 Patients should have easy access to reliable sources of information (web based and written)  
664 for joint replacement surgery. Options for anaesthesia and analgesia, and potential benefits  
665 and risks of each option should be discussed with patients to support shared decision  
666 making.<sup>65,125</sup>

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

667 9.2 Patient information leaflets on hip fractures should be available for patients, relatives and  
668 carers.<sup>126,127</sup> The clinicians should collaborate with the patients in selecting the most suitable  
669 treatment option based on best available evidence and informed patient  
670 preferences.<sup>67,128,129,130,131</sup>

671 9.3 Enhanced recovery programmes for patients undergoing primary arthroplasty surgery should  
672 provide comprehensive details of the patient journey including MDT led hip and knee school  
673 and expectations in terms of postoperative physiotherapy and early mobilisation where  
674 necessary. Information provided should be comprehensive and include details of regional  
675 anaesthesia.

676 9.4 When it is considered appropriate for a do not to attempt cardiopulmonary resuscitation in  
677 the event of a cardiopulmonary arrest (DNACPR) order, this should be discussed with  
678 capacitous patients, including those who have expressed their own wish not to be  
679 resuscitated.<sup>132</sup> In patients not without capacity to consent, every attempt should be made  
680 to discuss this with next of kin and/or patient advocates holding power of attorney (or an  
681 independent mental capacity advocate), according to local trust guidelines.

### 682 Consent

683 In order to give valid informed consent, patients need to understand the nature and purpose of the  
684 procedure. Detailed recommendations for consent are described in [GPAS Chapter 2: Guidelines  
685 for the Provision of Anaesthesia Services for the Perioperative Care of Elective and Urgent Care  
686 Patients](#).

687 9.5 Informed consent may not be possible for many patients undergoing hip fracture and major  
688 trauma surgery, owing to delirium, dementia, altered conscious level, severe pain or the  
689 effects of sedative drugs. Patients should not be asked to sign a consent form if they do not  
690 have capacity to do so. Standard operating procedures must be compliant with the Mental  
691 Capacity Act 2005. A high level of integrity should be maintained, and good documentation  
692 is essential.<sup>133,134,135</sup>

693 9.6 All decisions concerning the consent process and treatment plans, including decisions about  
694 whether or not to operate, should be documented clearly, noting the risks, benefits and  
695 alternatives that were explained to the patient.<sup>97,136</sup>

### 696 10 Financial considerations

697 Part of the methodology used in this chapter in making recommendations that considers the  
698 financial impact for each of the recommendations. Very few of the literature sources from which  
699 these recommendations have been drawn have included financial analysis.

700 The vast majority of the recommendations are not new recommendations, but they are a synthesis  
701 of already existing recommendations. The current compliance rates with many of the  
702 recommendations are unknown, and so it is not possible to calculate the financial impact of the  
703 recommendations in this chapter. It is impossible to make an overall assessment of the financial  
704 impact of these recommendations with the currently available information.

### 705 11 Implementation support

706 The Anaesthesia Clinical Services Accreditation (ACSA) scheme, run by the RCoA, aims to provide  
707 support for departments of anaesthesia to implement the recommendations contained in the  
708 GPAS chapters. The scheme provides a set of standards, and requires departments of anaesthesia  
709 to benchmark themselves against these using a self-assessment form available on the RCoA  
710 website. Every standard in ACSA is based on recommendation(s) contained in GPAS. The ACSA  
711 standards are reviewed annually and republished approximately four months following GPAS

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

712 review and republication, to ensure that they reflect current GPAS recommendations. ACSA  
713 standards include links to the relevant GPAS recommendations, for departments to refer to while  
714 working through their gap analyses.

715 Departments of anaesthesia are given the opportunity to engage with the ACSA process for an  
716 appropriate fee. Once engaged, departments are provided with a 'college guide', either a  
717 member of the ACSA committee or an experienced reviewer, to assist them with identifying actions  
718 required to meet the standards outlined in the document. Departments must demonstrate  
719 adherence to all 'priority one' standards listed in the document to receive accreditation from the  
720 RCoA. This is confirmed during a visit to the department by a group of four ACSA reviewers (two  
721 clinical reviewers, a lay reviewer and an administrator), who submit a report back to the ACSA  
722 committee.

723 The ACSA committee has committed to building a 'good practice library' (GPL), which will be used  
724 to collect and share documentation such as policies and checklists, as well as case studies of how  
725 departments that have overcome barriers to implementation of the standards, or have  
726 implemented the standards in innovative ways.

727 One of the outcomes of the ACSA process is to test the standards, and by extension the GPAS  
728 recommendations, to ensure that they are able to be implemented by departments of anaesthesia  
729 and consider any difficulties that may result from implementation. The ACSA committee has  
730 committed to measuring and reporting feedback of this type from departments engaging in the  
731 scheme back to the CDGs updating the guidance via the GPAS technical team.

### 732 Areas for future development

733 Following the systematic review of the evidence, the following areas of research are suggested:

- 734 • advanced pain relief techniques for trauma patients before surgery.

### 735 Abbreviations

ACSA	Anaesthesia Clinical Services Accreditation
CDG	Chapter Development Group
DAS	Difficult Airway Society
DNACPR	Do not to attempt resuscitation in the event of a cardiopulmonary arrest
ED	Emergency department
GMC	General Medical Council
GPAS	Guidelines for the Provision of Anaesthetic Services
LEH	Local emergency hospital
MDT	Multidisciplinary team
MRI	Magnetic resonance imaging
MTC	Major trauma centre
NICE	National Institute for Health and Care Excellence
NSCID	National Spinal Cord Injury Database
NSCIS	NHS Spinal Cord Injury Service
POCUS	Point of care ultrasound
RCoA	Royal College of Anaesthetists
SAS	Staff grade, associate specialist and specialty
TARN	Trauma Audit and Research Network
TU	Trauma unit
WHO	World Health Organization

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### 736 Glossary

737 **Autonomously practising anaesthetist** – a consultant or a staff grade, associate specialist or  
738 specialty (SAS) doctor who can function autonomously to a level of defined competencies, as  
739 agreed within local clinical governance frameworks.

740 **Clinical lead** – doctors undertaking lead roles should be autonomously practicing doctors who  
741 have competence, experience and communication skills in the specialist area equivalent to  
742 consultant colleagues. They should usually have experience in teaching and education relevant to  
743 the role and they should participate in Quality Improvement and CPD activities. Individuals should  
744 be fully supported by their Clinical Director and be provided with adequate time and resources to  
745 allow them to effectively undertake the lead role.

746 **Prehospital care** – auditing long term outcomes on fractured neck of femur and revision of major  
747 joint surgeries using a validated objective tool.

748 **Triage positive** – identified as severe injuries by the ambulance team using prehospital triage  
749 system.

750 **Triage negative** – not identified as severe injuries by the ambulance team using prehospital triage  
751 system.

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### References

- 1 The Trauma Audit and Research Network ([www.tarn.ac.uk](http://www.tarn.ac.uk))
- 2 Gleeson T, Blehar D. Point-of-care ultrasound in trauma. *Seminars in Ultrasound, CT & MRI* 2018; 39: 374-83
- 3 National Institute for Health and Care Excellence. Hypothermia: prevention and management in adults having surgery. NICE, 2016 ([bit.ly/3C7MTVK](https://bit.ly/3C7MTVK))
- 4 Royal College of Anaesthetists. Supervision of SAS anaesthetists in NHS hospitals. London, 2019 ([bit.ly/2THO1uo](https://bit.ly/2THO1uo))
- 5 Cook T, Woodall N, Frerk C. NAP4 Major Complications of Airway Management in the United Kingdom: 4th National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Report and findings. London: Royal College of Anaesthetists and Difficult Airway Society; 2011 ([bit.ly/34ildO5](https://bit.ly/34ildO5))
- 6 National Information Governance Board for Health and Social Care. Best practice tariff (BPT) for fragility hip fracture care user guide. The National Hip Fracture Database 2010 ([bit.ly/3EiXa47](https://bit.ly/3EiXa47))
- 7 Mallett S, Armstrong M. Point-of-care monitoring of haemostasis. *Anaesth* 2015; 70: 73-e26
- 8 World Health Organization. Guidelines for trauma quality improvement programmes. WHO, 2009 ([bit.ly/3yewf5G](https://bit.ly/3yewf5G))
- 9 Royal College of Anaesthetists. Chapter 11: Guidance for the Provision of Anaesthesia Services for Acute Inpatient Pain Management. RCoA, 2022 ([bit.ly/3M95C7Q](https://bit.ly/3M95C7Q))
- 10 Jansen J. Damage control resuscitation for patients with major trauma. *BMJ* 2009; 5: 338
- 11 Association of Anaesthetists. Regional analgesia for lower leg trauma and the risk of acute compartment syndrome. Association of Anaesthetists 2021 ([bit.ly/3SDX6jH](https://bit.ly/3SDX6jH))
- 12 NHS England. National Safety Standards for Invasive Procedures (NatSSIPs). London, 2015 ([bit.ly/1K6fRY2](https://bit.ly/1K6fRY2))
- 13 Health Improvement Scotland. Scottish Patient Safety Programme (SPSP). HIS ([bit.ly/3CtS8zh](https://bit.ly/3CtS8zh))
- 14 Regional Anaesthesia UK, Safe Anaesthesia Liaison Group. Stop Before you Block Campaign. London, 2015 ([bit.ly/2OhwwAl](https://bit.ly/2OhwwAl))
- 15 French J, Bedford N, Townsley P. Stop before you block campaign ([bit.ly/1IJYalm](https://bit.ly/1IJYalm))
- 16 National Institute for Health and Care Excellence. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism NICE guideline [NG89], March 2018 ([bit.ly/2pth9Wy](https://bit.ly/2pth9Wy))
- 17 BOAST: The care of the older or frail orthopaedic trauma patient. British orthopaedic association standard 2019 ([bit.ly/3SQSLcC](https://bit.ly/3SQSLcC))
- 18 Royal College of Anaesthetists. Chapter 2: Guidelines for the Provision of Anaesthesia Services for the Perioperative Care of Elective and Urgent Care Patients. RCoA, 2022 ([bit.ly/3V6wWrx](https://bit.ly/3V6wWrx))
- 19 Association Regional Analgesia for lower limb trauma: (<https://anaesthetists.org/Home/Resources-publications/Guidelines/Regional-analgesia-for-lower-leg-trauma-and-the-risk-of-acute-compartment-syndrome>)
- 20 National Institute for Health and Care Excellence. Major trauma: service delivery. NICE guideline [NG40]. NICE, 2016 ([bit.ly/3CddTCW](https://bit.ly/3CddTCW))
- 21 National Institute for Health and Care Excellence. Trauma Quality standards. NICE, 2018 ([bit.ly/2EjTX1u](https://bit.ly/2EjTX1u))
- 22 Kerr D, Bradshaw L, Kelly A-M. Implementation of the Canadian C-spine rule reduces cervical spine x-ray rate for alert patients with potential neck injury. *Journal of Emergency Medicine*. 2005;28: 127-31
- 23 Soar J, Deakin C, Lockey A, Nolan J, Perkins G. *Guidelines: Adult advanced life support*. London: Resuscitation Council UK; 2015 ([bit.ly/1MrJtKq](https://bit.ly/1MrJtKq))
- 24 British Orthopaedic Association. BOAST: The management of blunt chest wall trauma. British Orthopaedic Association Audit Standards for Trauma. BOA, 2017 ([bit.ly/3ygtzzM](https://bit.ly/3ygtzzM))
- 25 National Institute for Health and Care Excellence. Spinal injury: assessment and initial management. NICE, 2016 ([bit.ly/2DqRC48](https://bit.ly/2DqRC48))
- 26 British Orthopaedic Association. BOAST - Cervical Spine Clearance in the Trauma Patient. BOAST, 2021 ([bit.ly/3RHP6gc](https://bit.ly/3RHP6gc))
- 27 Royal College of Radiologists. Standards of practice and guidance for trauma radiology in severely injured patients. RCR, 2011 ([bit.ly/2FwUBdV](https://bit.ly/2FwUBdV))

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 28 Ramsingh D, Mangunta VR. The Use of Point-of-Care Ultrasonography in Trauma Anesthesia. *Anesthesiology Clinics* 2019; 37: 93-106
- 29 Royal College of Emergency Medicine. Handover; skills to enhancing the PHEM – EM interface. RCEM, 2021 ([bit.ly/3rAukEJ](https://bit.ly/3rAukEJ))
- 30 Müller M, Jürgens J, Redaelli M *et al.* Impact of the communication and patient hand-off tool SBAR on patient safety: a systematic review. *BMJ Open*, 2018; 8: 8
- 31 British Orthopaedic Association. BOAST - The Management of Traumatic Spinal Cord Injury. BOAST, 2014 ([bit.ly/3C7dRwI](https://bit.ly/3C7dRwI))
- 32 Behrens V, Dudaryk R, Nedeff N, Tobin JM, Varon AJ. The Ryder cognitive aid checklist for trauma. *Anesth Analg* 2016; 122: 1484-7
- 33 Faculty of Pre-Hospital Care. Pre-hospital spinal immobilisation: an initial consensus statement. *FPHC*, 2012 ([bit.ly/1YkqLQr](https://bit.ly/1YkqLQr))
- 34 Association of Anaesthetists of Great Britain and Ireland. *Recommendations for Standards of Monitoring During Anaesthesia and Recovery*, London 2015 ([bit.ly/1XVtRw](https://bit.ly/1XVtRw))
- 35 Association of Anaesthetists. AAGBI Safety Guideline. Immediate post-anaesthesia recovery 2013. London, 2013 ([bit.ly/30PJSI2](https://bit.ly/30PJSI2))
- 36 Gleeson T, Blehar D. Point-of-care ultrasound in trauma. *Seminars in Ultrasound, CT & MRI* 2018; 39: 374-83
- 37 Li L, Yong R, Kaye A, Urman R. Perioperative point of care ultrasound (POCUS) for anesthesiologists: an overview. *Current Pain and Headache Reports* 2020; 24: no 20
- 38 Ramsingh D, Mangunta V. The Use of Point-of-Care Ultrasonography in Trauma Anesthesia. *Anesthesiology Clinics* 2019; 37: 93-106
- 39 Multidisciplinary Association for Spinal Cord Injury Professionals. Management of people with spinal cord injury. *National Spinal Cord Injury Strategy Board*, 2011 ([bit.ly/1YkqSeJ](https://bit.ly/1YkqSeJ))
- 40 Spinal Injury Association ([www.spinal.co.uk](http://www.spinal.co.uk))
- 41 The Royal College of Radiologists. Standards of practice and guidance for trauma radiology in severely injured patients, 2011 ([bit.ly/3EfGIBy](https://bit.ly/3EfGIBy))
- 42 NHS standard contract for major trauma service. *NHSE*, 2013 ([bit.ly/2D27j6H](https://bit.ly/2D27j6H))
- 43 Advanced trauma and life support: student course manual. *Am Coll Surg*, 2009
- 44 Wathen JE, Gao D, Merritt G, Georgopoulos G, Battan FK. A randomized controlled trial comparing a fascia iliaca compartment nerve block to a traditional systemic analgesic for femur fractures in a pediatric emergency department. *Ann Emerg Med* 2007;50: 162-71
- 45 National Confidential Enquiry into Patient Outcome and Death. Trauma: who cares? A report of the National Confidential Enquiry into Patient Outcome and Death. *NCEPOD*, 2007 ([bit.ly/3M667PY](https://bit.ly/3M667PY))
- 46 Royal College of Anaesthetists. Chapter 5: Guidelines for the Provision of Emergency Anaesthesia. *RCoA*, 2022 ([bit.ly/3T1PuHz](https://bit.ly/3T1PuHz))
- 47 Association of Anaesthetists. Safe transfer of the brain-injured patient: trauma and stroke. Association of Anaesthetists, 2019 ([bit.ly/3EjRcQn](https://bit.ly/3EjRcQn))
- 48 Association of Anaesthetists. Management of Hip Fractures. Association of Anaesthetists, 2020 ([bit.ly/3CvzvP](https://bit.ly/3CvzvP))
- 49 National Institute for Health and Care Excellence. Joint replacement (primary): hip, knee and shoulder. *NICE*, 2020 ([bit.ly/3M7pc4d](https://bit.ly/3M7pc4d))
- 50 Hannon, Fillingham Y, Nam D *et al.* Opioids in Total Joint Arthroplasty: The Clinical Practice Guidelines of the American Association of Hip and Knee Surgeons, American Society of Regional Anesthesia and Pain Medicine, American Academy of Orthopaedic Surgeons, Hip Society, and Knee Society. *Journal of Arthroplasty* 2020; 35: 2709-2714a
- 51 Melamed R, Boland L, Normington J *et al.* A Postoperative respiratory failure necessitating transfer to the intensive care unit in orthopedic surgery patients: Risk factors, costs, and outcomes. *Perioperative Medicine* 2016; 5: 19
- 52 Jones J, Southerland W, Catalani B. The importance of optimizing acute pain in the orthopedic trauma patient. *Orthopedic Clinics of North America* 2017; 48: 445-65
- 53 Risk prediction in surgery [www.riskprediction.org.uk/index-nhfs.php](http://www.riskprediction.org.uk/index-nhfs.php)

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 54 Management of proximal femoral fractures 2011. *Anaesth* 2012; 67: 85-98
- 55 Facilities for surgical procedures: Volume 1. (HBN 26) DH, 2004 ([bit.ly/3SEB00s](https://bit.ly/3SEB00s))
- 56 NHS standard contract for major trauma service. *NHSE*, 2013 ([bit.ly/2D27j6H](https://bit.ly/2D27j6H))
- 57 Galvagno S, Nahmias J, Young D. Advanced trauma life support update 2019: management and applications for adults and special populations. *Anesthesiology Clinics* 2019; 37: 13-32
- 58 Sub-specialty training in pre-hospital emergency medicine. *IBTPEM*, 2012 ([bit.ly/2gPrW9g](https://bit.ly/2gPrW9g))
- 59 Statutory Instrument No.50: Blood Safety and Quality Regulations 2005. *HMSO*, 2005 ([bit.ly/23nrWP1](https://bit.ly/23nrWP1))
- 60 Griffiths R, Babu S, Dixon P *et al.* Guideline for the management of hip fractures. *Anaesthesia*, 2020; 76: 225-37
- 61 Royal College of Radiologists. Standards of practice and guidance for trauma radiology in severely injured patients. *RCR*, 2011 ([bit.ly/2FwUBdV](https://bit.ly/2FwUBdV))
- 62 Griffiths R, Surrendra Kuamar D, Major trauma in older people: implications for anaesthesia and intensive care medicine. *Anaesthesia* 2017; 72: 1302
- 63 Kamal T, Conway R, Littlejohn I, Ricketts D. The role of a multidisciplinary pre-assessment clinic in reducing mortality after complex orthopaedic surgery. *Ann R Coll Surg Engl* 2011; 93: 149-51
- 63 French J, Bedford N, Townsley P. Stop before you block campaign ([bit.ly/1JYalm](https://bit.ly/1JYalm))
- 64 Chan S, Ip K, Irwin M. Perioperative optimisation of elderly and frail patients a narrative review. *Anaesthesia* 2019; 74: 80-8
- 65 National Institute for Health and Care Excellence. NICE Joint replacement (primary): hip, knee and shoulder. NICE 2020 ([bit.ly/3CyZBxW](https://bit.ly/3CyZBxW))
- 66 Centre for Perioperative Care. Preassessment and optimisation for adult surgery. 2021 ([bit.ly/3Cdv33k](https://bit.ly/3Cdv33k))
- 67 Centre for Perioperative Care. Sharing decision making for Clinicians ([bit.ly/3CyW5DL](https://bit.ly/3CyW5DL))
- 68 National Institute for Health and Care Excellence. Shared Decision making. NICE 2021 ([bit.ly/3SFuC95](https://bit.ly/3SFuC95))
- 69 Henning J, Woods K. Management of major trauma. *Anaesth Intensive Care* 2014; 15: 405-7
- 70 Centre for Perioperative Care. CPOC Perioperative care of people living with frailty ([bit.ly/3T0tVqI](https://bit.ly/3T0tVqI))
- 71 Faculty of Pain Medicine *et al.* Surgery and Opioids: Best Practice Guidelines 2021. *FPM*, 2021 ([bit.ly/3yiUWOj](https://bit.ly/3yiUWOj))
- 72 Thomas W Wainwright, Mike Gill, David A McDonald, Robert G Middleton, Mike Reed, Opinder Sahota, Piers Yates & Olle Ljungqvist (2020) Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations, *Acta Orthopaedica*, 91:1, 3-19, DOI: [10.1080/17453674.2019.1683790](https://doi.org/10.1080/17453674.2019.1683790)
- 73 Hannon C, Fillingham Y, Nam D *et al.* Opioids in Total Joint Arthroplasty: The Clinical Practice Guidelines of the American Association of Hip and Knee Surgeons, American Society of Regional Anesthesia and Pain Medicine, American Academy of Orthopaedic Surgeons, Hip Society, and Knee Society. *Journal of Arthroplasty* 2020; 35: 2709-14
- 74 Melamed R, Boland L, Normington J *et al.* A Postoperative respiratory failure necessitating transfer to the intensive care unit in orthopedic surgery patients: Risk factors, costs, and outcomes. *Perioperative Medicine* 2016; 5: 19
- 75 Boloel H, Slim K. Sustainability of anaesthesia components of an enhanced recovery program (ERP) in colorectal and orthopaedics surgery. *Anaesthesia Critical Care & Pain Medicine* 2019; 38: 25-8
- 76 Gleicher Y, Singer O, Choi S, McHardy P. Thoracic epidural placement in a pre-operative block area improves operating room efficiency and decreases epidural failure rate. *Reg Anesth Pain Med* 2017; 42: 649-51
- 77 Perlman R, Callum J, Laflamme C *et al.* A recommended early goal-directed management guideline for the prevention of hypothermia-related transfusion, morbidity, and mortality in severely injured trauma patients. *Critical Care* 2016; 20: 107
- 78 Association of Anaesthetists. Cell salvage for peri-operative blood conservation. 2018 ([bit.ly/2mdU5MF](https://bit.ly/2mdU5MF))
- 79 Samolyk K, Beckmann S, Bissinger R. A new practical technique to reduce allogenic blood exposure and hospital costs while preserving clotting factors after cardiopulmonary bypass: the Hemobag. *Perfusion* 2005; 20: 343-9

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 80 Klein A, Arnold R, Bingham K *et al.* The use of blood components and their alternatives. *Anaesth* 2016; 71: 829-42
- 81 Walrath B, Harper S, Barnard E *et al.* Airway management for trauma patients. *Military Medicine* 2018; 183: 29-31
- 82 Setting up a difficult airway trolley (DAT). DAS ([bit.ly/1nJbXu0](https://bit.ly/1nJbXu0))
- 83 Bozzio A, Gala R, Villasenor M, Hao J, Mauffrey C. Orthopedic trauma surgery in the morbidly obese patient. *Eur J Orthop Surg Traumatol* 2014; 24: 421-5
- 84 Galvagno S, Nahmias J, Young D. Advanced trauma life support update 2019: management and applications for adults and special populations. *Anesthesiology Clinics* 2019; 37: 13-32
- 85 Mallett S, Armstrong M. Point-of-care monitoring of haemostasis. *Anaesth* 2015; 70: 73-e26
- 86 Brazzel C. Thromboelastography-guided transfusion Therapy in the trauma patient. *AANA Journal* 2013; 81: 127-32
- 87 Statutory Instrument No.50: Blood Safety and Quality Regulations 2005. HMSO, 2005 ([bit.ly/23nrWP1](https://bit.ly/23nrWP1))
- 88 Facilities for surgical procedures: Volume 1. (HBN 26) DH, 2004 ([bit.ly/1RtwFu0](https://bit.ly/1RtwFu0))
- 89 CRASH-2 collaborators. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet* 2010; 376: 23-32
- 90 CRASH-2 collaborators. The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomised controlled trial. *Lancet* 2011; 377: 1096-101
- 91 Fillingham Y, Ramkumar D, Jevsevar D *et al.* Tranexamic Acid use in total joint arthroplasty: the clinical practice guidelines endorsed by the American Association of Hip & Knee Surgeons, American Society of Regional Anesthesia and Pain Medicine, American Academy of Orthopaedic Surgeons, Hip Society and Knee Society. *Journal of Athroplasty* 2018; 33: 3065-9
- 92 Kietaiabl A, Kietaiabl S. New hemostatic agents: Perioperative anesthetic considerations. *Current Pharmaceutical Design*. 2019; 25: 2158-64
- 93 General Medical Council. Protecting children and young people: the responsibilities of all doctors. GMC, 2012 ([bit.ly/1cNoXZk](https://bit.ly/1cNoXZk))
- 94 The Children Act 2004. HMSO, 2004 ([bit.ly/1snJHsF](https://bit.ly/1snJHsF))
- 95 MacArthur B, Foley M, Gray K, Sisley A. Trauma in Pregnancy: A Comprehensive Approach to the Mother and Fetus. *American Journal of Obstetrics & Gynecology* 2019; 220: 465-8
- 96 Cheek TG, Baird E. Anesthesia for nonobstetric surgery: maternal and fetal considerations. *Clin Obs Gyne* 2009; 52: 535-45
- 97 Upadya M, Sanest P. Anaesthesia for non-obstetric surgery during pregnancy. *Indian J Anaesth* 2016; 60: 234-41
- 98 Adams M, Nomellini V, Bennett S. Anesthetic concerns in the pregnant trauma patient. *Current Trauma Reports* 2018; 4: 218-24
- 99 Hill K, Gross M, Sutton K, Mulcahey M. Evaluation and resuscitation of the pregnant orthopaedic trauma patient: considerations for maternal and fetal outcomes. *JBS Reviews* 2019; 7: e3
- 100 The Trauma Audit and Resarch Network. Major Trauma in Older People. TARN, 2017 ([bit.ly/3Mb1Jil](https://bit.ly/3Mb1Jil))
- 101 Fisher J, Bates C, Banerjee J. The growing challenge of major trauma in older people: a role for comprehensive geriatric assessment? *Age & Ageing* 2017; 46: 709-12
- 102 Fort A, Zack-Guasp R. Anesthesia for patients with extensive trauma. *Anesthesiology Clinics* 2020; 38: 135-148
- 103 Oakley P, Dawes R, Rhys Thomas GO. The consultant in trauma resuscitation and anaesthesia. *BJA* 2014; 113: 207-10
- 104 Royal College of Anaesthetists. 2021 *Anaesthetics Curriculum*, 2021 ([bit.ly/34DcQMz](https://bit.ly/34DcQMz))
- 105 Sub-specialty training in pre-hospital emergency medicine. IBPEM, 2012 ([bit.ly/2gPrW9g](https://bit.ly/2gPrW9g))
- 106 Fort A, Zack-Guasp R. Anesthesia for patients with extensive trauma. *Anesthesiology Clinics* 2020; 38: 135-48
- 107 Ditzel R, Anderson J, Eisenhart W *et al.* A review of transfusion and trauma induced hypocalcemia: Is it time to change the lethal triad to the lethal diamond? *J Trauma Acute Care Surg*. 2020; 88: 434-9

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- 108 Wray J, Bridwell R, Schauer S *et al*. The diamond of death: Hypocalcemia in trauma and resuscitation. *Am J Emerg Med*. 2021; 41: 104-9
- 109 Schmitz B, Zern S. Training for trauma anesthesia: Role of education and simulation based training. *Current Anesthesiology Reports* 2020; 10: 196-203
- 110 Hutchins D, Nightingal T, Pollard L, Rockett M. Chest trauma pathway and effects of regional anaesthesia at a major trauma centre. *British Journal of Pain* 2018; 12 (2 Suppl 1): 39-40
- 111 Pouryahya P, Meyer A, Koo M, Chu T. The utility of point-of-care ultrasound (POCUS) in emergency department; an observational study. *Emergency Medicine Australasia* 2018; 30 (suppl 1) 6-7
- 112 Li L, Yong R, Kaye A, Urman R. Perioperative point of care ultrasound (POCUS) for anesthesiologists: an overview. *Current Pain and Headache Reports* 2020; 24: no 20
- 113 Association of Anaesthetists. Implementing advance care plans in the peri-operative period, including plans for cardiopulmonary resuscitation. Association of Anaesthetists, 2022 ([bit.ly/3TOK0NK](https://bit.ly/3TOK0NK))
- 114 Beaudoin F, Haran J, Liebmann O. A comparison of ultrasound-guided three-in-one femoral nerve block versus parenteral opioids alone for analgesia in emergency department patients with hip fractures: a randomized controlled trial. *Acad Emerg Med* 2013; 20: 584-91
- 115 Wathen JE, Gao D, Merritt G, Georgopoulos G, Battan FK. A randomized controlled trial comparing a fascia iliaca compartment nerve block to a traditional systemic analgesic for femur fractures in a pediatric emergency department. *Ann Emerg Med* 2007;50: 162-7
- 116 National Institute for Health and Care Research. Trauma and Emergency Care ([bit.ly/3fLoCNB](https://bit.ly/3fLoCNB))
- 117 National Institute for Health and Care Excellence. Major trauma: assessment and initial management. NICE, 2018 ([bit.ly/3ecbVuQ](https://bit.ly/3ecbVuQ))
- 118 Home office. Policy paper: Saving and Improving Lives: The Future of UK Clinical Research Delivery, 2021 ([bit.ly/3e691HR](https://bit.ly/3e691HR))
- 119 The National Hip Fracture Database. RCP, London ([www.nhfd.co.uk](http://www.nhfd.co.uk))
- 120 National Joint Registry ([www.njrcentre.org.uk](http://www.njrcentre.org.uk))
- 121 National Peer Review Programme: major trauma measures (version 1.0, November 2014). NHSE ([bit.ly/2o8zaf1](https://bit.ly/2o8zaf1))
- 122 Elkassabany N, Cai L, Mehta S *et al*. Does regional anesthesia improve the quality of postoperative pain management and the quality of recovery in patients undergoing operative repair of Tibia and Ankle fractures? *J Orthop Trauma*. 2015; 29: 404-9
- 123 Moro E, Silva M, Couri M, Issa D, Barbieri J. Quality of recovery from anesthesia in patients undergoing orthopedic surgery of the lower limbs. *Brazilian Journal of Anesthesiology* 2016; 66: 642-50
- 124 Ramsingh D, Mangunta V. The Use of Point-of-Care Ultrasonography in Trauma Anesthesia. *Anesthesiology Clinics* 2019; 37: 93-106
- 125 Groves ND, Humphreys H, Williams A, Jones A. Effect of informational internet web pages on patients' decision-making: randomised controlled trial regarding choice of spinal or general anaesthesia for orthopaedic surgery. *Anaesth* 2010; 65: 277-82
- 126 National Institute for Health and Care Excellence. Hip fracture: management. Clinical guideline [CG124]. NICE, 2017 ([bit.ly/3SUP7OX](https://bit.ly/3SUP7OX))
- 127 Royal College of Anaesthetists and Association of Anaesthetists. Anaesthetic choices for hip or knee replacement. RCoA, 2020 ([bit.ly/3gicqnS](https://bit.ly/3gicqnS))
- 128 Royal College of Anaesthetists, Association of Anaesthetists and Regional Anaesthesia UK. Nerve blocks for surgery on the shoulder, arm or hand. RCoA, 2020 ([bit.ly/3VM8Hzc](https://bit.ly/3VM8Hzc))
- 129 Royal College of Anaesthetists. Preparing for a knee arthroscopy ([bit.ly/3eGAdxq](https://bit.ly/3eGAdxq))
- 130 Royal College of Anaesthetists. Preparing for a knee replacement operation ([bit.ly/3gioeGr](https://bit.ly/3gioeGr))
- 131 Royal College of Anaesthetists, Association of Anaesthetists and Regional Anaesthesia UK. Your spinal anaesthetic. RCoA, 2020 ([bit.ly/3ENADfs](https://bit.ly/3ENADfs))
- 132 Meek, T, Clyburn, R, Fritz, Z, Pitcher, D, Ruck Keene A, Young P. Implementing advance care plans in the peri-operative period, including plans for cardiopulmonary resuscitation: Association of Anaesthetists clinical practice guideline. *Anaesthesia*, 2022; 77: 456-62
- 133 Mental Capacity Act 2005. HMSO, 2005 ([bit.ly/1Hz3HDZ](https://bit.ly/1Hz3HDZ))

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

134 Adult With Incapacity (Scotland) Act 2000. HMSO, 2000 ([bit.ly/1ntNhpO](http://bit.ly/1ntNhpO))

135 Mental Capacity Act (Northern Ireland) 2016 ([bit.ly/2wDApVr](http://bit.ly/2wDApVr))

136 The Children Act 2004. HMSO, 2004 ([bit.ly/1snJHsF](http://bit.ly/1snJHsF))

DRAFT

### Appendix 1: Recommendations grading

The grading system is outlined in the methodology section of this chapter. The grades for each of the recommendations in this chapter are detailed in the table below:

TBC

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### About these guidelines

#### Methodology

The process by which this chapter has been developed has been documented within the GPAS Chapter Development Process Document, which is available on request.

The evidence included in this chapter is based on a systematic search of the literature. Abstracts were independently screened by two investigators and reviewed against inclusion and exclusion criteria. Data were extracted by one investigator in accordance with predefined criteria. The review objective was to determine the key components needed to ensure provision of high-quality perioperative services for patients who have undergone surgery and/or interventions which involve anaesthesia.

#### Search strategy

Searches were performed on **TBC**, for the literature search strategy, outcomes, databases, criteria for inclusion and exclusion of evidence (for the full Neuroanaesthetic services chapter search protocol please contact the RCoA). A hand search of the literature was also conducted by the authors using the reference lists of relevant original articles and review articles.

The literature search was performed in **TBC**.

The authors and researcher independently reviewed the abstracts and titles of the studies found in the initial search. After agreement on the primary selection of papers, full-text versions were accessed and reviewed against the following predefined inclusion and exclusion criteria. The full-text papers were also reviewed by the CDG for suitability. The final list of publications used can be found in the references.

#### Inclusion criteria

The literature review considered studies that included the following patient population with all of the inclusion criteria listed below:

- all patients undergoing elective or emergency anaesthesia
- all staff groups working within Neuroanaesthetic care, under the responsibility of an anaesthetic clinical director, including (but not restricted to) consultant anaesthetists, autonomously practising anaesthetists, anaesthetists in training, nurses, operating department practitioners, surgeons, pharmacists, general practitioners, radiologists and radiographers.

#### Exclusion criteria

The literature review used the following exclusion criteria:

- provision of neuroanaesthesia provided by a speciality other than anaesthesia.

#### Data extraction and analysis

Data were extracted by the authors using a proforma. The study characteristics data included:

- the journal and country of publication
- the number of patients recruited into the study
- the study design
- patient characteristics
- outcome data

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

- the logic of the argument
- author's conclusions
- reviewer's comments.

The patient characteristics data extracted were: age, gender and type of surgery. The analysis considers studies that included any clinical outcome, including (but not restricted to) survival, length of stay – critical care or hospital, morbidity, adverse effects and complications.

The evidence that is included in this chapter has been graded according to a grading system adapted from NICE and outlined below:

Level	Type of evidence	Grade	Evidence
<b>Ia</b>	Evidence obtained from a single large/multicentre randomised controlled trial, a meta-analysis of randomised controlled trials or a systematic review with a low risk of bias	<b>A</b>	At least one randomised controlled trial as part of a body of literature of overall good quality and consistency addressing the specific recommendation (evidence level I) without extrapolation
<b>Ib</b>	Evidence obtained from meta-analyses, systematic reviews of RCTs or RCTs with a high risk of bias	<b>B</b>	Well-conducted clinical studies but no high-quality randomised clinical trials on the topic of recommendation (evidence levels Ib, II or III); or extrapolated from level Ia evidence
<b>IIa</b>	Evidence obtained from at least one well-designed controlled study without randomisation		
<b>IIb</b>	Evidence obtained from at least one well-designed quasi-experimental study		
<b>IIc</b>	Evidence obtained from case control or cohort studies with a high risk of confounding bias		
<b>III</b>	Evidence obtained from well-designed non-experimental descriptive studies, such as comparative studies, correlation studies and case studies		
<b>IV</b>	Evidence obtained from expert committee reports or opinions and/or clinical experiences of respected authorities	<b>C</b>	Expert committee reports or opinions and/or clinical experiences of respected authorities (evidence level IV) or extrapolated from level I or II evidence. This grading indicates that directly applicable clinical studies of good quality are absent or not readily available.
<b>UG</b>	Legislative or statutory requirements	<b>M</b>	This grading indicates that implementation of this recommendation

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

			is a statutory requirement, or is required by a regulatory body (e.g. CQC, GMC)
		<b>GPP</b>	Recommended good practice based on the clinical experience of the CDG.
<p><b>Adapted from Eccles M, Mason J. How to develop cost-conscious guidelines. <i>Health Technology Assessment</i> 2001;5(16) and Mann T. Clinical guidelines: using clinical guidelines to improve patient care within the NHS. <i>Department of Health</i>, London 1996.</b></p>			

### Strengths and limitations of body of evidence

Most of the published evidence on perioperative care anaesthesia services is descriptive. There are publications describing aspects of this process based on expert opinion.

The limitations of the evidence are:

- the 'unmeasurables' (attitudes, behaviour, motivation, leadership, teamwork)
- few randomised controlled trials (RCTs); studies frequently use mixed populations of emergency and elective patients, or all emergency patients grouped together despite different underlying diagnoses
- papers often examine a single intervention within complex system or bundle
- papers are often examining small numbers and/or patients from a single centre
- poor use of outcome measures, frequently concentrating on easily measured short-term outcomes which are not patient centred
- generally, a paucity of long-term follow up
- there is no standard definition used of 'high risk'
- use of different risk-scoring systems
- decrease in outcome over time and geography when 'good papers' are used in quality improvement programmes
- application of international studies in systems with either more or less resources than the UK into NHS practice
- older studies may no longer be applicable within the NHS
- very few studies included any analysis of financial implications
- evidence was mainly based on literature graded III and IV.

### Methods used to arrive at recommendations

Recommendations were initially drafted based on the evidence by the authors for the chapter. These were discussed with the CDG, and comments were received both on the content and the practicality of the recommendations. The level of evidence that was the basis for each recommendation was graded according to a grading system, and the recommendation was then graded taking into account the strength of the evidence and the clinical importance using a recommendations criteria form.

Recommendations were worded using the following system of categorisation:

Strength	Type of evidence	Wording
----------	------------------	---------

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

<b>Mandatory</b>	The evidence supporting the recommendation includes at least one with an 'M' grading	Wording should reflect the mandatory nature of the recommendation i.e. 'must'
<b>Strong</b>	Confidence that for the vast majority of people, the action will do more good than harm (or more harm than good)	Wording should be clearly directive 'should' or 'should not'
<b>Weak</b>	The action will do more good than harm for most patients, but may include caveats on the quality or size of evidence base or patient preferences	Wording should include 'should be considered'
<b>Aspirational</b>	While there is some evidence that implementation of the recommendation could improve patient care, either the evidence or the improvement is not proven or substantial	Wording should include 'could'
<b>Equipose</b>	There is no current evidence on this recommendation's effect on patient care	Wording should include 'there is no evidence of this recommendation's effect on patient care'

### Consultation

The chapter has undergone several rounds of consultation. The multidisciplinary CDG formed the first part of the consultation process. The authors and GPAS Editorial board identified key stakeholder groups. Where stakeholders are represented by an association or other medical college, they were asked to nominate delegates to join the CDG. The GPAS Chapter Development Process Document (available on request) explains the recruitment process for those CDG members who were not directly nominated. The CDG members were involved in drafting the recommendations, and were provided with an opportunity to comment on all subsequent drafts of the chapter.

The chapter underwent peer review. Peer reviewers were identified by the GPAS Editorial Board, Clinical Quality and Research Board (CQRB) or through the Clinical Leaders in Anaesthesia Network. Nominees were either anaesthetists of consultant grade or were nominated by a key stakeholder group. Nominees had not had any involvement in the development of GPAS to date and were asked to comment upon a late draft of the chapter.

Following peer review, the chapter was reviewed by the College's CQRB and the College's Lay Committee. Comments from all groups were considered and incorporated into a consultation draft.

The consultation draft of this chapter was circulated for public consultation from TBC. As well as being made available on the College's website and promoted via Twitter and the President's newsletter to members, the draft was also circulated to all key stakeholder groups identified by the authors and the College. A list of organisations contacted by the College is available from the GPAS team at the College: [GPAS@rcoa.ac.uk](mailto:GPAS@rcoa.ac.uk).

# Chapter 16

## Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

### The editorial independence of GPAS

The development of GPAS is wholly funded by the Royal College of Anaesthetists. However, only the GPAS technical team and the GPAS researcher are paid directly by the College for their work on GPAS: the GPAS Editors' employing organisation receives 2 programmed activities (PA) backfill funding. All funding decisions by the College are made by the chief executive officer, in collaboration with the senior management team and College Council.

The authors of the chapters are all fellows of the Royal College of Anaesthetists. Members of College Council cannot act as chair of any CDG, as this individual has the deciding vote under the consensus method of decision making used in the chapters. Where College Council members have been involved in chapter development, this has been declared and recorded.

All persons involved in the development of GPAS are required to declare any pecuniary or non-pecuniary conflict of interest, in line with the GPAS conflict of interest policy as described in the GPAS Chapter Development Process Document (available on request). Any conflicts of interest are managed on a case-by-case basis to maintain the transparency and impartiality of the GPAS document. The conflicts, and the way they were managed, are outlined at the beginning of the chapter.

### The role of the GPAS Editorial Board and CQRB

The overall development of the entire GPAS document is overseen by the CQRB of the Royal College of Anaesthetists, which includes representatives from all grades of anaesthetist and from clinical directors, and which also has lay representation.

Responsibility for managing the scope of the document and providing clinical oversight to the project technical team is delegated by the CQRB to the GPAS Editorial Board, which includes individuals responsible for the various internal stakeholders (see above for membership). On the inclusion/exclusion of specific recommendations within each chapter, the Editorial Board can only provide advice to the authors. In the event of disagreement between the authors, the majority rules consensus method is used, with the GPAS Editor holding the deciding vote.

Both of these groups, along with the College's Lay Committee, review each chapter and provide comment prior to public consultation and are responsible for signoff before final publication. In the event of disagreement, consensus is reached using the majority rules consensus method, with the chair of CQRB holding the deciding vote.

### Updating these guidelines

This chapter will be updated for republication in January 2025.

Guidelines will be updated on an annual basis. The researcher will conduct the literature search again using the same search strategy to uncover any new evidence and members of the public will be able to submit new evidence to the GPAS project team. Where new evidence is uncovered, the lead author will decide whether the recommendations that were originally made are still valid in light of this new evidence.

If new evidence contradicts or strengthens existing recommendations, the authors decide whether or not to involve the remainder of the CDG in revising the recommendations accordingly.

If new evidence agrees with existing recommendations, then a reference may be added but no further action is required.

If there is no new evidence then no action is required.

This chapter is due to be fully reviewed for publication in January 2028.

## Chapter 16

# Guidelines for the Provision of Anaesthesia Services for Trauma and Orthopaedic Surgery 2023

Every five years guidance will be submitted to a full review involving reconvening the CDG (or appointment of a new, appropriately qualified CDG), and the process described in the methodology section of this chapter begins again.



Royal College of Anaesthetists

Royal College of Anaesthetists, Churchill House, 35 Red Lion Square, London WC1R 4SG  
020 7092 1500 | [www.rcoa.ac.uk/gpas](http://www.rcoa.ac.uk/gpas) | [GPAS@rcoa.ac.uk](mailto:GPAS@rcoa.ac.uk)

© Royal College of Anaesthetists (RCoA)

DRAFT