

References

- 1 Boyle A. Unprecedented? The NHS crisis in emergency care was entirely predictable. *BMJ* 2023;Jan 9;380:46. doi: 10.1136/bmj.p46. PMID: 36623878.
- 2 Quince W. 2023 March 30. [Hospital Beds: Costs.](#)
- 3 The high-risk general surgical patient: raising the standard. [RCS Eng, 2018.](#)
- 4 Eugene N, Oliver CM, Bassett MG, Poulton TE, Kuryba A, Johnston C, Anderson ID, Moonesinghe SR, Grocott MP, Murray DM, Cromwell DA, Walker K; NELA collaboration. Development and internal validation of a novel risk adjustment model for adult patients undergoing emergency laparotomy surgery: the National Emergency Laparotomy Audit risk model. *Br J Anaesth* 2018;Oct;121(4):739–748. doi: 10.1016/j.bja.2018.06.026. Epub 2018 Aug 2023. PMID: 30236236.
- 5 NELA Project Team. Eighth Patient Report of the National Emergency Laparotomy Audit. [RCOA, London 2023.](#)
- 6 NELA Project Team. Ninth Patient Report of the National Emergency Laparotomy Audit. [RCOA, London 2024.](#)
- 7 Bion J, Barton G, Boyle A, Carlson G, Carrol E, Christian W, Crossland S, Faust S, Gabbay F, Gent N, Gibbs P, Harden J, Howard P, Inada-Kim M, Lanzman M, Leanord A, Lees N, Noursadeghi M, Rajesh P, Shrestha S, Sriskandan S, Stockley S, Waldmann C, Walsh T, Wilson P, Singer M. Academy of Medical Royal Colleges Statement on the Initial Antimicrobial Treatment of Sepsis. *AoMRC*, 2022.
- 8 Wong DJN, Popham S, Wilson AM, Barneto LM, Lindsay HA, Farmer L, Saunders D, Wallace S, Campbell D, Myles PS, Harris SK, Moonesinghe SR; SNAP-2: EPICCS collaborators; Study Steering Group. Postoperative critical care and high-acuity care provision in the United Kingdom, Australia, and New Zealand. *Br J Anaesth* 2019;Apr 122(4):460–469. doi: 10.1016/j.bja.2018.12.026. Epub 2019 Feb 8. PMID: 30857602; PMCID: PMC6435907.