

Table 5.1: Characteristics of NoLap patients

	Number (n)	Percentage (%)
Total patients analysed	943	100
Age group		
18 - 29	<10	<1.0
30 - 39	<10	<1.0
40 - 49	14	1.5
50 - 59	48	5.1
60 - 69	96	10.2
70 - 79	237	25.1
80 - 89	363	38.5
90 +	178	18.9
Patients aged 65+	837	88.8
Sex		
Male	390	41.4
Female	553	58.6
Risk status*		
High-risk	924	98
Low risk	19	2
Bowel pathology		
Bowel perforation	459	48.7
Bowel ischaemia	330	35.0
Bowel perforation and ischaemia	58	6.2
Not stated	96	10.2
Type of admission		
Elective	11	1.2
Emergency	932	98.8
Admitting specialty		
General surgery	662	70.2
General medicine	192	20.4

	Number (n)	Percentage (%)
Total patients analysed	943	100
Age group		
18 - 29	<10	<1.0
30 - 39	<10	<1.0
40 - 49	14	1.5
50 - 59	48	5.1
60 - 69	96	10.2
70 - 79	237	25.1
80 - 89	363	38.5
Other	41	4.3
Elderly Care	26	2.8
Oncology	14	1.5
Gastroenterology	<10	<1.0
Gynaecology (including Gynae-Oncology)	<10	<1.0

*Risk status is defined by either a NELA Parsimonious risk score (PRS) of 5% or greater, patients living with frailty- defined by a Rockwood Clinical Frailty Scale (CFS) of 5 or more Clinical Frailty Scale (CFS) or indicated from clinical assessment based on other conditions not included in PRS.