

Nerve blocks for surgery on the shoulder, arm or hand


Information for patients and families



Nerve blocks for surgery on the shoulder, arm or hand

This leaflet is for anyone who is thinking about having a nerve block for an operation on the shoulder, arm or hand. It will be of particular interest to people who would prefer not to have a general anaesthetic.

The leaflet has been written with the help of patients who have had a nerve block for their operation.



Brachial plexus block?

The brachial plexus is the group of nerves that lies between your neck and your armpit. It contains all the nerves that supply movement and feeling to your arm – from your shoulder to your fingertips.

A brachial plexus block is an injection of local anaesthetic around the brachial plexus. It 'blocks' information travelling along these nerves. It is a type of nerve block. Your arm becomes numb and immobile.

You can then have your operation without feeling anything. The block can also provide excellent pain relief for between three and 24 hours, depending on what kind of local anaesthetic is used.

A brachial plexus block rarely affects the rest of the body so it is particularly advantageous for patients who have medical conditions which put them at a higher risk for a general anaesthetic.

A brachial plexus block may be combined with a general anaesthetic or with sedation. This means you have the advantage of the pain relief provided by a brachial plexus block, but you are also unconscious or sedated during the operation.

Your anaesthetist can explain what is possible and what might be best for you.

Other nerve blocks

If your operation is on the hand or forearm, it may be possible to do a nerve block on nerves further down the arm, closer to the hand. Your anaesthetist can tell you more.

Having the injection

You need to prepare for the operation by not eating or drinking. The hospital should give you clear instructions about this.


Nerve blocks for surgery on the shoulder, arm or hand

You will go to a room near the operating theatre to have the local anaesthetic injection.

The injection for a brachial plexus block is in the side of your neck, or in your armpit, or close to your collar bone. Other nerves can be blocked near the elbow, or in the forearm, wrist or hand.

Sedation is commonly given before the injection is done. This is when drugs are given which help you relax. If you are having a general anaesthetic as well, this may be given before or after the injection.

The skin around the injection site is cleaned. A small injection of local anaesthetic numbs the skin. The nerves are located using an ultrasound machine or by using a small machine that makes your arm twitch.



Most people find that the injection is no more painful than having a cannula inserted into a vein. Please see the leaflet [You and your anaesthetic](http://www.rcoa.ac.uk/patientinfo/leaflets-video-resources) on our website for more information about cannulas: www.rcoa.ac.uk/patientinfo/leaflets-video-resources

Your arm will start to feel warm, heavy and numb. The injection takes between 20 and 40 minutes to work.

Sometimes the block does not work fully. This may be due to the operation being more extensive than expected or due to technical difficulty with the injection. If this happens, you will be offered more local anaesthetic, additional pain relief, or a general anaesthetic.

During the operation

A screen is used, so you cannot see the operation being done, unless you want to.

A member of staff will sit with you during the operation. Your anaesthetist remains close by. Please feel free to bring in a personal music player with headphones if you would like to listen to music during the operation.

An operating theatre is a busy place – there will usually be between five and eight people in theatre, each with their own role in helping look after you.

Nerve blocks for surgery on the shoulder, arm or hand

If you are having sedation, you will be relaxed and drowsy. You may be given oxygen through a light plastic facemask. You may have some memories of being in the operating theatre, although these will probably be patchy.

If you have a general anaesthetic you will not remember anything about being in the operating theatre.

Benefits of a brachial plexus block

Better pain relief afterwards. There may be less need for strong pain relief medicines which make some people feel quite sick and unwell.

Avoiding a general anaesthetic, including its risks and side effects. The common side effects of a general anaesthetic include sickness, sore throat and drowsiness.






Often able to leave hospital sooner.

Side-effects, complications and risks

In modern anaesthesia, serious problems are uncommon. Risk cannot be removed completely, but modern drugs, equipment and training have made anaesthesia a much safer procedure in recent years.

Anaesthetists take a lot of care to avoid all the risks given in this section. Your anaesthetist will be able to give you more information about any of these risks and the precautions taken to avoid them. You can also find out more information from the patient information pages on the College website: www.rcoa.ac.uk/patientinfo.

People vary in how they interpret words and numbers. This scale is provided to help.

				
Very common	Common	Uncommon	Rare	Very rare
1 in 10	1 in 100	1 in 1,000	1 in 10,000	1 in 100,000
One person in your family	One person in a street	One person in a village	One person in a small town	One person in a large town

- **Injection in the side of the neck:** hoarse voice, droopy eyelid, some difficulty breathing. These resolve as the block wears off.
- **Injection around the collar bone:** less than 1 in 1,000 risk of damage to the covering of the lung. Your anaesthetist will discuss this with you.
- **All injection sites:** damage to a blood vessel which usually resolves with simple compression to stop any bleeding.
- **Very rarely:** having a fit or another life threatening event may occur. Your anaesthetist will manage these promptly. They can tell you more about these very rare events.

Nerve blocks for surgery on the shoulder, arm or hand

Nerve damage

- The risk of long-term nerve damage caused by a brachial plexus block is difficult to measure precisely. Studies show that it happens in between 1 in 700 and 1 in 5,000 blocks.
- If you have a block lower down the arm, the risk of nerve damage and its consequences may be less.
- About 1 in 10 patients notice a prolonged patch of numbness or tingling in their arm. These symptoms will resolve in 95% of these patients within four to six weeks, and in 99% within a year.
- There is a risk of nerve damage after any operation regardless of whether you have had a block. This can be due to the operation, the position you lie in or the use of a tourniquet (tight band on the upper arm which prevents bleeding during the operation). Swelling around the operation site or a pre-existing medical condition, such as diabetes, may also contribute to nerve damage.

For more information on nerve damage, please see the leaflet Nerve damage associated with peripheral nerve block on our website: www.rcoa.ac.uk/patientinfo/risks/risk-leaflets

After the operation

During the time the block is working you will not be able to use your arm. You will probably be given a sling and you may need someone to help you look after yourself.

You should start taking pain relief medicines while your arm is still numb and before the block wears off. This is so that they start working ready for when the block wears off. When this happens you may experience pins and needles in your fingers – this is normal.

Aftercare

You should keep your arm in the sling you are given, for support and protection. You will not be fully aware of the position of your arm – so it can be injured without you realising.

- Be especially careful around heat sources, such as fires or radiators. You will not feel heat while your arm is numb and burns can happen.
- Avoid use of any machinery or domestic appliances. Injury is possible while you cannot feel your arm.
- Start taking your pain relief medicines before the block wears off. This is important as the pain can appear quite suddenly.

Further help

You will need to seek help from the emergency medical services if you notice unexplained breathlessness, or severe pain that is not controlled by your tablets.

Nerve blocks for surgery on the shoulder, arm or hand

If the block has not fully worn off by 48–72 hours after the operation you should contact your anaesthetist by calling the anaesthetic department at your hospital.

Questions you may like to ask your anaesthetist

- 1 Who will be doing the injection?
- 2 Who will stay with me during the operation?
- 3 What happens if the block does not work and I can feel something? How often does this happen?
- 4 Do I have any particular risks for having this kind of anaesthetic?
- 5 Do I have increased risk for a general anaesthetic, which I might have instead of a brachial plexus block?
- 6 When will my arm feel normal again?
- 7 Who should I call if I am concerned about the after effects of the block?

You can find more information leaflets on the College website www.rcoa.ac.uk/patientinfo. The leaflets may also be available from the anaesthetic department or pre-assessment clinic in your hospital.

Risks associated with your anaesthetic

Leaflets about specific risks associated with having an anaesthetic or an anaesthetic procedure are also available via the College website: www.rcoa.ac.uk/patientinfo/risks/risk-leaflets. They supplement the patient information leaflets listed above.

Tell us what you think

We welcome suggestions to improve this leaflet. If you have any comments that you would like to make, please email them to patientinformation@rcoa.ac.uk

Royal College of Anaesthetists

Churchill House, 35 Red Lion Square, London WC1R 4SG

020 7092 1500 | patientinformation@rcoa.ac.uk | www.rcoa.ac.uk/patientinfo

 @RCoANews

 RoyalCollegeofAnaesthetists

First Edition 2015

This leaflet will be reviewed within five years of the date of publication.

© Royal College of Anaesthetists (RCoA)

This leaflet may be copied for the purpose of producing patient information materials. Please quote the original source.

If you wish to use part of this leaflet in another publication, suitable acknowledgement must be given and the logos, images and branding removed. For more information, please contact us.